PREVENTIVE EFFECT OF DIETARY *CUCURBITA MAXIMA* DUCH FRUIT INCLUSION ON N- METHYL-N-NITROSOUREA (MNU) INDUCED COLON CARCINOGENESIS IN MALE WISTAR RATS

By

IBRAHIM ISIAKA

DEPARTMENT OF BIOCHEMISTRY
FACULTY OF LIFE SCIENCES
AHMADU BELLO UNIVERSITY, ZARIA
NIGERIA.

DECEMBER 2017

PREVENTIVE EFFECT OF DIETARY *CUCURBITA MAXIMA* FRUIT DUCH INCLUSION ON N- METHYL-N-NITROSOUREA (MNU) INDUCED COLON CARCINOGENESIS IN MALE WISTAR RATS

By

Ibrahim ISIAKA

(P14SCBC8093)

A THESIS SUBMITTED TO THE SCHOOL OF POSTGRADUATE STUDIES, AHMADU BELLO UNIVERSITY, ZARIA

IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE AWARD OF MASTERS DEGREE IN NUTRITION.

DEPARTMENT OF BIOCHEMISTRY
FACULTY OF LIFE SCIENCES
AHMADU BELLO UNIVERSITY, ZARIA
NIGERIA

DECEMBER 2017

DECLARATION

I declare that the work in this Dissertation entitled **"Preventive Effect of Dietary *Cucurbita maxima* Fruit (Duch) Inclusion on N-Methyl-N-Nitrosourea (MNU) Induced Colon Carcinogenesis in Wistar Rats"** has been performed by me in the Department of Biochemistry. The information derived from the literature has been duly acknowledged in the text and a list of references provided. No part of this Dissertation was previously presented for another degree or diploma at this or any other Institution.

IBRAHIM ISIAKA	Signature	Date

CERTIFICATION

This Dissertation entitled **"Preventive Effect of Dietary *Cucurbita maxima* Fruit (Duch) Inclusion on N- Methyl-N-Nitrosourea (MNU) Induced Colon Carcinogenesis in Wistar Rats"** by **IBRAHIM ISIAKA** meets the regulations governing the award of the degree of Master of Science in Nutrition of the Ahmadu Bello University, and is approved for its contribution to knowledge and literary presentation.

Prof. S.E. Atawodi, FAS

Chairman, Supervisory Committee _____ Signature _____ Date

Prof. K. M. Anigo

Member, Supervisory Committee _____ Signature _____ Date

Prof. M. N Shuaibu

Head of Department _____ Signature _____ Date

Prof. S. Z Abubakar

Dean, School of Postgraduate Studies _____ Signature _____ Date

ACKNOWLEDGEMENTS

My sincere gratitude goes to Almighty God for being my only source of strength, hope, helper and provider in every phase of my studies. Alhamdulillah Allah, I can't thank you enough.

I sincerely appreciate my Supervisor, guardian and my confidant, Professor S.E. Atawodi FAS, for all the support, professional advice and immense contributions towards my success in this race. May you live long and forever be in total sound health Sir. My gratitude also to my Co-supervisor Professor K.M. Anigo for his contribution.

I thank the Head of Biochemistry Department for permission to use available facilities. I sincerely would love to thank my academic father and mentor, Professor D.A Ameh for all the knowledge impacted in me and the recommendations. I thank Dufil Prima Foods, for the 2015 Postgraduate Scholarship in Nutrition.

To my mother and everything, Mrs. Hauwa'u Isiaka, I can't thank you enough for your prayers both day and night to see that I succeed in everything. I love you mother and may our good God whom we serve continue to keep you for me, Ameen.

I would not forget to show my sincere appreciation to my colleagues in this renowned research work (the cancer research group, a.k.a My MNU family), Dr. Mubarak Liman, Abdulrazaq Yusuf, Oguche Alhassan Solomon, Late John Ofodile, Itanyi Job Onuche and Jamila Lawal. You all are too kind and I pray that God continues to favor us all in our life endeavors. Ameen! Ameeen!!

I thank my siblings for their encouragement, prayers and supports; Hajiya Maryam Mustapha, Hassanah Isiaka, Muhammed Kabir Isiaka, Dr. Rabbi Isiaka, Accountant Suleiman Isiaka, Engineer Isah Isiaka, Nurse Ameenah Isiaka Audi, and my best friend Oladele Omolade Victor, without you all, this work would not have been possible. Thank you very much and God bless you all.

ABSTRACT

The preventive effect of dietary C*ucurbita maxima* fruit inclusion at 2.5%, 5% and 10% on colon carcinogenesis initiated with N-methyl-N-nitrosourea (MNU) induction for twelve (12) weeks was evaluated. Group A= Normal saline with standard feed, Group B= MNU with standard feed, Group C= MNU with dietary inclusion of 2.5% *Cucurbita maxima* powder, Group D= MNU with dietary inclusion of 5% *Cucurbita maxima powder*, Group E= MNU with dietary inclusion of 10% *Cucurbita maxima* powder, Group F= Dietary inclusion control normal saline with 10% Cucurbita maxima powder. After twelve (12) weeks of treatment, the colon and liver of rats undergoing MNU intoxication but fed *Cucurbita maxima* fruit included diets showed a significant ($p \leq 0.05$) increase in the activity of endogenous antioxidant enzymes (catalase and superoxide dismutase) when compared with MNU control group. Also a dose-dependent decrease in the extent of lipid peroxidation was observed in the colon and liver of dietary *Cucurbita maxima* fruit included treated groups. A significant ($p \leq 0.05$) increase in white blood cell, platelet level, neutrophils and lymphocyte counts in MNU control group was observed when compared to other groups fed with dietary *Cucurbita maxima* fruit inclusion, while a significant ($p \leq 0.05$) decrease in packed cell volume, haemoglobin count and red blood cell were observed in MNU control group compared to other groups fed with dietary *Cucurbia maxima* fruit inclusion. Histological observation of colon and liver tissue sections showed severe focal area of inflammatory cells in the MNU control group while normal control group and other groups that received various dietary *Cucurbita maxima* fruit inclusion showed milder to no tissue damage. A statistically significant ($p \leq 0.05$) decrease in carcino-embrayonic antigen (CEA) levels was observsed in all dietary *Cucurbita maxima* fruit inclusion groups compared to MNU control group. Immunohistochemical observations of the colon tissue sections showed no expression of

Mut L Homolog gene in MNU control group while moderate to mild expression of *Mut L Homolog* gene in normal control group and other groups treated with 2.5%, 5% and 10% of dietary *Cucurbita maxima* fruit inclusion. These results indicate potential of dietary *Cucurbita maxima* fruit inclusion to antagonizing the process of chemically induced colon carcinogenesis by enhancement of endogenous antioxidants systems and prevention of oxidative stress.

TABLE OF CONTENTS

Title page.....i
Declaration.....ii
Certification.....iii
Dedication.....iv
Acknowledgements.....v
Table of Contents.....vi
List of Figures.....x
List of Tables.....xi
List of Plates.....xii
List of Appendices.....xiii
List of Abbreviations.....xiv
Abstract.....xvi
CHAPTER ONE**.....1**
1.0 *INTRODUCTION*.....*1*
1.1 **Statement of Researc Problem**.....3
1.2 **Justification**.....4
1.3 **General Aim**.....5
1.4 **Specific Objectives**.....5
1.5 **Null Hypothesis**.....5
CHAPTER TWO.....6
2.0 **LITERATURE REVIEW**.....6
2.1 **Colon**.....6
2.2 **Colon Cancer**.....7
2.3 **Incidence of Colon Cancer**.....7

2.4 **Causes and Mechanism of Colon Carcinogenesis…….....**9

2.5 Some Diagnostic (Biomarkers) in Colon Carcinogenesis……14

2.5.1 Carcinoembryonic Antigen……..…....14

2.5.2 Normal range……..……14

2.5.3 **Sensitivity……**15

2.5.4 Monitoring treatment……....15

2.5.8 Carcinoembryonic Antigen as a marker for colorectal cancer screening……17

2.6 Immunohistochemistry……..18

2.6.1 Basic Principles of Immunohistochemistry…….....19

2.6.2 Principles of Immunohistochemistry…….....20

2.7 Animal Model in Colon Carcinogenesis…….....21

2.7.1 Rodent Colon Tumor Models (History and Overview)……..21

2.8 **Role of MNU in Colon Carcinogenesis…….....**23

2.9 Haematological Parameters in Cancer/Colon Carcinogenesis and their Application as Possible Cancer Biomarker…….....24

2.9.1 Haematological Components and their Functions……..25

2.10 Oxidative Stress and Cancer / Colon Carcinogenesis…….....27

2.11 Antioxidant……..28

2.12 *Cucurbita Species*……....29

2.12.1 Description of plant……29

2.12.2 Taxonomical Position of *Cucurbita maximus*……29

2.13 Origin and Distribution…….....32

2.14 Culinary Uses…….....33

2.15 Medicinal Uses…….....33

2.16 **Chemical Composition……...**34

2.17 Known Pharmacological Activities…….....35

2.17.1 Anticancer Activity- Anticancer activity of methanol extract of *Cucurbita maxima* Against Ehrlich as-cites carcinoma……...35

2.17.2 Antiobesity Activity of Cucurbita Maxima Seed Extract With Beta Vulgaris & Smilax Regelii Root Extract To Reduce Extra Fats from the Body……...36

2.17.3 Antidiabetic Activity – Antidiabetic activity of *Cucurbita maxima* aerial parts…....36

2.17.4 Hepatoprotective Activity - Evaluation of Hepatoprotective Activity of *Cucurbita maxima* Aerial Parts……....37

2.17.5 Diuretic Activity - Diuretic activity of seeds of *Cucurbita maxima* duchesne in albino wistar rats…….37

CHAPTER THREE……...40

3.0 MATERIALS AND METHODS……..40

3.1 Chemicals……..40

3.2 Apparatus and Equipment…….....40

3.3 Sample Collection and Preparation…….....41

3.4 Experimental Diets……...41

3.5 Experimental Animals……...41

3.6 Induction of Colon Cancer……...44

3.7 Collection of Organs or Tissues……...44

3.8 Estimation of Endogenous Antioxidants…….....45

3.8.1 Lipid peroxidation by measuring the malondialdehyde (MDA) level……45

3.8.2 Superoxide Dismutase (SOD) activity……...45

3.8.3 Catalase assay……46

3.9 Histological Analysis of the Colon and other Organs……47

3.10 Haematological Analysis……47

3.11 Carcinoembryonic Antigen Assay…….....47

3.12 Immunohistochemical Analysis…….....48

3.13 Statistical Analysis……..49

CHAPTER FOUR……50

4.0 RESULTS…….50

CHAPTER FIVE……..64

5.0 RESULT AND DISCUSSION…….....64

CHAPTER SIX……..69

6.0 SUMMARY, CONCLUSION AND RECOMMENDATIONS…….....69

6.1 SUMMARY……...69

6.2 CONCLUSION……..69

6.3 RECOMMENDATION……...69

REFERENCES……...70

APPENDICES……81

LIST OF FIGURES

Figure 2.1: Anatomy of the large intestine showing the four sections: caecum and ascending, transverse, descending and sigmoid. The rectum forms the most distal portion of the large intestine……..7

Figure: 2.3: Source: International Journal of Phytopharmacy……...31

Figure 3.1: EXPERIMENTAL DESIGN for the Dietary Inclusion *Cucurbita maxima* Fruit for Prevention of Colon Carcinogenesis in Male Albino Wistar Rats…….....43

Figure 4.1: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Packed Cell Volume, Neutrophil and Lymphocyte Values following 12 Weeks MNU Administration in different Experimental Groups……..51

Figure 4.2: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Haemoglobin and Platelet Concentration following 12 Weeks MNU Administration in Different Experimental Groups…….....52

Figure 4.3: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on White Blood Cell and Red Blood Cell following 12 Weeks MNU Administration in Different Experimental Groups…….....54

Figure 7: CEA Concentration of the Serum following Intrarectal Administration of MNU in Rats Fed with different Level of Dietary *C. maxim* Fruit Inclusion…….....62

LIST OF TABLES

Table 4.1: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Superoxide dismutase Activity following 12 Weeks Intrarectal MNU Instillation.……56

Table 4.2: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Catalase Activity following 12 Weeks Intrarectal MNU Instillation ……...57

Table 4.3: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Superoxide dismutase Activity following 12 Weeks Intrarectal MNU Instillation …….58

LIST OF PLATES

Plate: 2.4: Fruit of different Sizes, Shapes and Color for *Cucurbita maxima* ……..…….31

Plate: 2.5: *Cucurbita maxima* Plantation Showing Leaves of Vegetable Plant, Flower and Fruit from Ochi Ogbonicha in Ofu LGA Kogi State, Nigeria……..39

Plates 4.1: Photomicrogragh of Histopathological Section of the Colon following 12 Weeks MNU Administration Fed with Different Level of Dietary *Cucurbita maxima* Fruit Inclusion in Rats…….....60

Plate 4.2: The Photomicrogragh of Histopathological Section of the Liver following 12 Weeks MNU Administration Fed with Different Level of Dietary *Cucurbita maxima* Fruit Inclusion in Rats……61

Plates 4.1: Photomicrogragh of Immunohistochemical Section of the Colon following 12 Weeks MNU Administration Fed with Different Level of Dietary *Cucurbita maxima* Fruit Inclusion in Rats…….....63

LIST OF APPENDICES

APPENDIX 1.0 Effect of *Cucurbita maxima* fruit on Percentage Packed Cell Volume (PCV), Neutrophil and Lymphocyte in the Blood following 12 Weeks MNU Instillation in Different Experimental Group…….....81

APPENDIX 2.0 Effect of *Cucurbita maxima* fruit on Haemoglobin and Platelet Concentration in the Blood Serum following 12 Weeks MNU Instillation in Different Experimental group…….....82

APPENDIX 3.0 Effect of *Cucurbita maxima* Fruit on White Blood Cell and Red Blood Cell in the Blood following 12 Weeks MNU Instillation in Different Experimental Group…….....83

APPENDIX 4.0 Effect of Dietary *Cucurbita maxima* Fruit Inclusion on MDA Level following 12 Weeks MNU Instillation in Different Experimental Group …….84

APPENDIX 5.0 Effect of Dietary *Cucurbita maxima* Fruit Inclusion of Catalase Activity following 12 Weeks MNU Instillation in Different Experimental Group……..85

APPENDIX 6.0 Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Superoxide Dismutase Activity following 12 Weeks MNU Instillation in Different Experimental Group……86

APPENDIX 7.0 CEA Concentration of the Serum following Intrarectal Administration of MNU in Rats Fed with Different Level of Dietary Inclusion of *C. maxima* Fruit…….....87

LIST OF ABBREVIATIONS AND SYMBOLS

ABC	Avidin Biotin Complex
ACF	Aberrant Crypt Foci
ANOVA	Analysis of Variance
AORTIC	African Organization for Research and Training in Cancer
APC	Adenomatous Polyposis Coli
AST	Aspartate Aminotransferase
BHA	Butylated Hydroxyl Anisole
BHT	Butylated Hydroxyl Toluene
CAT	Catalase
CEA	Carcinoembrayonic antigen
CIN	Chromosomal Instability
CRC	Colorectal Cancer
DPPH	1, 1-Diphenyl-2-picrylhydrazyl
DNA	Deoxyribonucleic acid
ELISA	Enzyme-linked immunosorbent assay
EVF	Erythrocyte Volume Fraction
GLOBOCAN	Global Burden on Cancer
GSHPx	Glutathione Peroxidase
Hb	Haemoglobin
HNE	Hydroxynonenal
Ht or Hct	Haematocrit

IBD	Inflammatory bowel diseases
MDA	Malondialdehyde
MSI	Microsatellite instability
MLH1	*Mult L homologue* 1
MMR	Mismatch Repair Gene
MNU	N-methyl nitrosourea
NADPH	Nicotinamide Adenine dinucleotide Phosphate Reduced
NCI	National Cancer Institute
PCV	Packed cell volume
PUFAs	Polyunsaturated Fatty Acids
RBC	Red blood cell count
RNS	Reactive nitrogen specie
ROS	Reactive Oxygen Species
SOD	Superoxide Dismutase
TAG	Triacylglyerol
TBA	Thiobarbituric Acid
WBC	White Blood Cell
WHO	World health organization

CHAPTER ONE

2.0 INTRODUCTION

Cancer is a group of diseases characterized by uncontrolled growth and spread of abnormal cells. If the spread is not controlled, it can result in death. Cancer is caused by both external factors (tobacco, chemicals, radiation, and infectious organisms) and internal factors (inherited mutations, hormones, immune conditions, and mutations that occur from metabolism). These causal factors may act together or in sequence to initiate or promote carcinogenesis. The development of most cancers requires multiple steps that occur over many years. Certain types of cancer can be prevented by eliminating exposure to tobacco and other factors that initiate or accelerate this process. Other potential malignancies can be detected before cells become cancerous or at an early stage, when the disease is most treatable. Cancer is treated with surgery, radiation, chemotherapy, hormones, and immunotherapy. Worldwide, one in eight deaths is due to cancer, cancer causes more deaths than AIDS, tuberculosis, and malaria combined. When countries are grouped according to economic development, cancer is the leading cause of death in developed countries and the second leading cause of death in developing countries (following heart diseases). According to recent World Health Organization (WHO) projections, cancer will have replaced ischemic heart disease as the overall leading cause of death worldwide in 2010 (WHO, 2007).

According to estimates from the International Agency for Research on Cancer (IARC), there were 12.7 million new cancer cases in 2008 worldwide, of which 5.6 million occurred in economically developed countries and 7.1 million in economically developing countries. The corresponding estimates for total cancer deaths in 2008 were 7.6 million (about 21,000 cancer deaths a day), 2.8 million in economically developed countries and 4.8 million in economically

developing countries. By 2030, the global burden is expected to grow to 21.4 million new cancer cases and 13.2 million cancer deaths simply due to the growth and aging of the population, as well as reductions in childhood mortality and deaths from infectious diseases in developing countries Ferlay *et al*., (2010).

The estimated future burden could be much larger than given above due to the adoption of western lifestyles, such as smoking, poor diet, physical inactivity, and reproductive factors, in economically developing countries. Cancers related to these factors, such as lung, breast, and colorectal cancers, are increasing in economically transitioning countries. Rates of cancers common in Western countries will continue to rise in developing countries if preventive measures are not widely applied. In economically developed countries, the three most commonly diagnosed cancers were prostate, lung and bronchus, and colorectal among men, and breast, colorectal, and lung among women. In economically developing countries, the three most commonly diagnosed cancers were lung, stomach, and liver in men, and breast, cervix uteri, and lung in women. In both economically developed and developing countries, the three most common cancer sites were also the three leading causes of cancer death.

Globally, colorectal cancer (CRC) is the second most common cancer in women (614, 000 cases per year) and the third most common in men (746, 000 cases per year). The incidence rates are higher in more developed countries (737, 000 cases per year) than in less developed ones (624, 000 cases per year). However, mortality is higher in the latter (52% of total deaths), which indicates poor survival. In 2015, the GLOBOCAN online analysis tool has predicted 61, 228 new CRC cases for Asia. Accordingly, 25, 816 of these cases are associated with people who are less than 65 years old Fearon *et al.,* (2011).

1.1 Statement of Research Problem

Cancer is still one of the most deadly diseases that cause serious health problems, physical disabilities, mortalities, and morbidities around the world. It is the second leading cause of death all over the world. Although great advancement has been made in the treatment of cancer progression, there is still significant deficiencies and room for improvement Sabira Sultana *et al.,* (2015). Colorectal carcinoma is the commonest malignancy of the gastrointestinal tract worldwide, with about 140,000 cases each year in the United States, according to the National Cancer Institute. Incidence and death rates for colorectal cancer increase with age. This means that about 90% of new cases and 93% of deaths occur in people 50 and older (SEER program 1969-2010). It is observed that the median age of colon cancer diagnosis is put at 69 in men and 73 in women, and that of rectal cancer diagnosis is put at 63 in men and 65 in women. Overall, colorectal cancer incidence and mortality rates are about 30% to 40% higher in men than in women (Murphy *et al.,* 2010). The reasons for this likely reflect complex interactions between gender-related differences in exposure to hormones and risk factors. However, the incidence is rising in Africans from an average of 10-20 newly diagnosed colorectal cancer in a tertiary health institution per year to about 50 Archampong *et al.,* (1994). Others are the complications that accompany resection and the damages caused by radiotherapy recently; resistance to anticancer drugs has been observed therefore, more research is required to explore the anticancer or preventive potential of dietary supplementation with medicinal plants (Vinay *et al.,* 2011).

1.2 Justification

Cancer is the second leading cause of death worldwide. Conventional cancer therapies cause serious side effects and at best, merely extend the patient's lifespan by a few years. The demand to utilize alternative concepts or approaches to the treatment of cancer is therefore escalating. However, the potential treatment of cancer is still under investigation. In fact the plants occupy a good place in the treatment Srijita-Dutta *et al.,* (2015).

The medicinal plants and their products, particularly vegetables have antioxidant activity leading to anticancer effect. Many doctors recommend that people who wish to reduce the risk of cancer must eat vegetables every day in their diet Srijita-Dutta *et al.,* (2015). The vegetables contain many phytochemicals having antioxidant activity. The antioxidant protects the cells from damage caused by free oxygen radicals. Here in this study, we will give a better therapeutic approach to cancer by the maximum use of vegetable (*Cucurbita maxima* fruit.) against colon cancer.

The fruit of *Cucurbita maxima* can be found in some part of Nigeria and it is used among the Hausa, igala and ibo ethnic tribes as day to day food and vegetables in various delicacies. Substantial evidence from various experimental studies has also suggested the anticancer potential of the different parts of the plant including the fruits on different types of cancer. Thus, considering the increasing incidence of colon cancer worldwide and the need for approaching and solving the problem of colon cancer through diet (dietary inclusion), there is therefore the need to investigate its potential (*Cucurbita spp*.) to prevent colon cancer Attarde *et al*., (2010).

1.3 General Aim

The aim of this work is to evaluate the preventive effect of dietary *Cucurbita maxima* fruit inclusion on MNU induced colon carcinogenesis in albino Wister rats.

1.4 Specific Objectives

i. To determine the effect of *Cucurbita maxima* fruit dietary inclusion on haematological parameters in MNU induced colon carcinogenesis in Albino wistar rats.

ii. To establish the possible role of *Cucurbita maxima* fruit included diet on the antioxidant status in MNU induced colon carcinogenesis in Albino wistar rats.

iii. To evaluate the effect of included diet of *Cucurbita maxima* fruit on some markers of MNU induced colon carcinogenesis in Albino wistar rats.

1.5 Null Hypothesis

Dietary *Cucurbita maxima* fruit inclusion has no preventive effect on colon carcinogenesis.


CHAPTER TWO

2.0 LITERATURE REVIEW

2.1 The Human Colon

The colon is an organ shaped like a question mark **(?)** that forms part of the large intestine (Figure1) (NCCN, 2014). It is approximately 1.5 metres long and is divided into four segments based on the vascular supply to each segment. The right colon, otherwise known as the ascending colon, receives its blood supply from the right colic and the ileocolic artery. The most proximal portion of the right colon is termed the caecum and is identified by the appendix, which a small appendage attached to three longitudinal bands of muscle called the taeniae coli. The transverse colon connects the right colon to the left colon. Its blood supply is derived from the middle colic artery and, like the right colic artery, it is derived from the superior mesenteric artery, a branch of the aorta. The juncture at which the right and transverse colon join is termed the hepatic flexure and the juncture at which the transverse colon and left colon join is termed the splenic flexure. Flexures are points at which the large bowel is tethered and supported in order to prevent excessive twisting that could lead to a condition called volvulus Mobarhan *et al.,* (2012). Volvulus is a serious problem that can result in diminished blood supply to the bowel, with subsequent necrosis and perforation. The transverse colon also serves as a tethering point for the greater omentum, which is a large fatty apron of tissue that essentially covers the entire abdomen, separating the small and large bowel from the anterior abdominal wall. The purpose of the omentum is to 'police' the abdomen, identifying holes or perforations of the bowel wall and sealing them. In addition, it seems to be an adhesive target for neoplastic cells,

which otherwise float free in the abdomen. The left colon connects the transverse colon to the sigmoid colon, which is named for its S-shaped appearance in the abdomen. The left colon and sigmoid colon are supplied by the left colic and sigmoidal arteries, branches of the inferior mesenteric artery Mobarhan *et al.*, (2012).



Source: Encyclopedia of Life Sciences 2001

FIG. 2.1: Anatomy of the large intestine showing the four sections: caecum and ascending, transverse, descending and sigmoid. The rectum forms the most distal portion of the large intestine.

2.2 Colon Cancer

According to Encyclopedia of life sciences 2001, Colon cancer is a disease of the large intestine which begins at a structure called the caecum, located in the right lower quadrant of the abdomen, and continues through all portions of the abdomen to its junction with the rectum, located in the deep pelvis.

2.3 Incidence of Colon Cancer

Colorectal cancer is the third most common cancer in men and the second in women Landis *et al.,* (*2011*). Worldwide, an estimated 1.2 million cases of colorectal cancer occurred in 2008. The highest incidence rates were in North America, Australia, New Zealand, Europe, and Japan. Rates were low in Africa and South Central Asia. Rates were substantially higher in men than in women, Cummings *et al.,* (*2011*).

The incidence of colorectal cancer is increasing in certain countries where risk was historically low (Japan), (Cummings *et al.,* 2011). In high-risk/high-income countries, trends are either gradually increasing (Finland, Norway), stabilizing (France, Australia), or declining (United States) with time. The greatest increases in the incidence of colorectal cancer are in Asia (Japan, Kuwait, Israel) and Eastern Europe (Czech Republic, Slovakia, Slovenia). In fact, rates among males in the Czech Republic, Slovakia, and Japan have not only exceeded the peak rates observed in long-standing developed countries such as the United States, Canada, and Australia but they continue to increase, Cummings (2011). The decrease in colorectal cancer incidence in the United States partially reflects the increase in detection and removal of precancerous lesions; the increase in several Asian and Eastern European countries may reflect changes in risk factors for colorectal cancer that are associated with westernization such as elevated obesity and smoking prevalence. In contrast to incidence trends, decreasing colorectal cancer mortality rates have been observed in a large number of countries worldwide and are most likely attributed to colorectal cancer screening and/or improved treatments (GLOBOCAN, 2008). However, increases in mortality rates are still occurring in countries that have more limited resources, including Mexico and Brazil in South America and Romania and Russia in Eastern Europe.

About 608,700 deaths from colorectal cancer occurred in 2008 worldwide, accounting for 8% of all cancer deaths, Cummings *et al.,* (2011).

Although inadequate research opportunities and data collection systems raise concerns of possible under-reporting, available data suggest a lower burden of CRC in Africa. While the overwhelming majority of the population in Africa is of indigenous African origin (black), countries with substantial other racial groups, such as in South Africa, report significant population (racial) differences in CRC burden. The lowest burden of disease have been noted among blacks, and especially those who have maintained their traditional lifestyle including diet, an intermediate burden among Indians and those of mixed ancestry (coloreds) and the highest burden among persons of European origin (whites). This contrasts the findings in the United States where African Americans suffer a disproportionally higher burden of CRC when compared to other population groups Cummings *et al.,* (2011).

Colorectal carcinoma (CRC) is relatively uncommon in Nigeria, compared to the developed countries. In 2008, colorectal cancer was the third commonest cancer in males and the second in females worldwide. However almost 60% of these cases occurred in developed countries and only 2% of new cases and 3% of deaths due to colorectal cancer occurred in Africa (Globocan 2008). Recent studies of CRC indicate an increasing trend, which has been attributed to the adoption of sedentary western lifestyle and diet, migration patterns and the improved availability of requisite specialized medical services Duduyemi *et al.,* (2013).

2.4 Causes and Mechanism of Colon Carcinogenesis

Carcinogenesis is the formation of a cancer, whereby normal cells are transformed into cancer cells. The process is characterized by changes at the cellular, genetic, and epigenetic levels and abnormal cell division, in some cancers forming a malignant mass Lim *et al.*, (2006). Cell division is a physiological process that occurs in almost all tissues and under many circumstances. Under normal circumstances, the balance between proliferation and programmed cell death, usually in the form of apoptosis, is maintained by regulation of both processes to ensure the integrity of tissues and organs. According to the prevailing accepted theory of carcinogenesis Tariq *et al.,* (2016), the somatic mutation theory, mutations and epimutations in DNA that lead to cancer (only certain mutations and epimutations can lead to cancer and the majority of potential mutations and epimutations will have no such effect) disrupt these orderly processes by disrupting the programming regulating the processes, upsetting the normal balance between proliferation and cell death. This results in uncontrolled cell division and the evolution of those cells by natural selection in the body. The uncontrolled and often rapid proliferation of cells can lead to benign tumors; some types of these may turn into malignant tumors (cancer). Benign tumors do not spread to other parts of the body or invade other tissues, and they are rarely a threat to life unless they compress vital structures or are physiologically active, for instance, producing a hormone. Malignant tumors can invade other organs, spread to distant locations (metastasis) and become life-threatening Tariq *et al.,* (2016).

More than one mutation is necessary for carcinogenesis. In fact, a series of several mutations to certain classes of genes is usually required before a normal cell will transform into a cancer cell Dow *et al.,* (2015). On average, for example, 15 “driver mutations” and 60 “passenger”

mutations are found in colon cancers, Wood *et al.,* (2007). Mutations in those certain types of genes that play vital roles in cell division, apoptosis (cell death), and mutations and epimutations in DNA repair genes will cause a cell to lose control of its cell proliferation, Weisenberger *et al.,* (2010).

Oncovirinae, viruses that contain an oncogene, are categorized as oncogenic because they trigger the growth of tumorous tissues in the host. This process is also referred to as viral transformation. Cancer is fundamentally a disease of regulation of tissue growth. In order for a normal cell to transform into a cancer cell, genes that regulate cell growth and differentiation must be altered, Crose (2008). Genetic and epigenetic changes can occur at many levels, from gain or loss of entire chromosomes, to a mutation affecting a single DNA nucleotide, or to silencing or activating microRNA that controls expression of 100 to 500 genes, Lim *et al.,* (2005)*;* Balaguer *et al.,* (2010). There are two broad categories of genes that are affected by these changes; Oncogenes may be normal genes that are expressed at inappropriately high levels, or altered genes that have novel properties. In either case, expression of these genes promotes the malignant phenotype of cancer cells. Tumor suppressor genes are genes that inhibit cell division, survival, or other properties of cancer cells. Tumor suppressor genes are often disabled by cancer promoting genetic changes. Typically, changes in many genes are required to transform a normal cell into a cancer cell, Fearon *et al.,* (2009).

Damage of DNA is considered to be the primary cause of cancer Bonadon, (2008); Bernstein *et al.,* (2011). More than 60,000 new naturally occurring DNA damages arise, on average, per human cell, per day, due to endogenous cellular processes. Additional DNA damages can arise from exposure to exogenous agents. As one example of an exogenous carcinogenic agent,

tobacco smoke causes increased DNA damage, and these DNA damages likely cause the increase of lung cancer due to smoking, Cunningham *et al.,* (2011). In other examples, UV light from solar radiation causes DNA damage that is important in melanoma, Przybylowska *et al.*, (2013) H. pylori infection produces high levels of reactive oxygen species that damage DNA and contributes to gastric cancer, Dow *et al.,* (2015), and the Aspergillus metabolite, aflatoxin, is a DNA damaging agent that is causative in liver cancer, Bernstein *et al.,* (2011). DNA damages can also be caused by endogenous (naturally occurring) agents. Katsurano *et al*., (2011) indicated that macrophages and neutrophils in an inflamed colonic epithelium are the source of reactive oxygen species causing the DNA damages that initiate colonic tumorigenesis, Katsurano *et al.*, (2011) and bile acids, at high levels in the colons of humans eating a high fat diet, also cause DNA damage and contribute to colon cancer Bernstein *et al.*, (2011).

A deficiency in DNA repair would cause more DNA damages to accumulate, and increase the risk for cancer. For example, individuals with an inherited impairment in any of 34 DNA repair genes are at increased risk of cancer with some defects causing up to 100% lifetime chance of cancer, Tariq *et al.*, (2016).

Colorectal cancer (CRCs) can arise from one or a combination of three different mechanisms, namely chromosomal instability (CIN), CpG island methylator phenotype (CIMP), and microsatellite instability (MSI). According to Ferlay *et al.*, (2015), the classical CIN pathway begins with the acquisition of mutations in the adenomatous polyposis coli (*APC*), followed by the mutational activation of oncogene *KRAS* and the inactivation of the tumor suppressor gene, *TP53*. Aneuploidy and loss of heterozygosity (LOH) are the major players in CIN tumors, which

not only constitute most of the sporadic tumors (85%) but also involve familial adenomatous polyposis cases associated with germline mutations in the *APC* gene. The CIMP pathway is characterized by promoter hypermethylation of various tumor suppressor genes, most importantly *MGMT* and *MLH1*. This hypermethylation is often associated with *BRAF* mutation and microsatellite instability Weisenberger *et al.,* (2010). The MSI pathway involves the inactivation of genetic alterations in short repeated sequences. This activation occurs in CRCs in DNA mismatch repair (MMR) genes, and is a hallmark condition in familial Lynch syndrome (LS), which also appears in ~15% of the sporadic CRC cases. In addition, the hypermethylation of the MMR genes may lead to MSI. This mechanism is often associated with the CIMP pathway Weisenberger *et al.,* (2010). MSI tumors are often associated with proximal colon and poor differentiation but better prognosis. Moreover, the three mechanisms often overlap in CRC.

In order for cells to start dividing uncontrollably, genes that regulate cell growth must be dysregulated Bernstein *et al.,* (2011). Proto oncogenes are genes that promote cell growth and mitosis, whereas tumor suppressor genes discourage cell growth, or temporarily halt cell division to carry out DNA repair. Typically, a series of several mutations to these genes is required before a normal cell transforms into a cancer cell. This concept is sometimes termed "oncoevolution." Mutations to these genes provide the signals for tumor cells to start dividing uncontrollably. But the uncontrolled cell division that characterizes cancer also requires that the dividing cell duplicates all its cellular components to create two daughter cells. The activation of anaerobic glycolysis (the Warburg effect), which is not necessarily induced by mutations in proto-oncogenes and tumor suppressor genes, Tariq *et al.,* (2016) provides most of the building blocks

required to duplicate the cellular components of a dividing cell and, therefore, is also essential for carcinogenesis Lopez *et al.*, (2010).

2.5 Some Diagnostic (Biomarkers) in Colon Carcinogenesis

2.5.1 Carcinoembryonic Antigen

Carcinoembryonic antigen (CEA) is a glycoprotein, which is present in normal mucosal cells but increased amounts are associated with adenocarcinoma, especially colorectal cancer. CEA therefore has a role as a tumor marker. Sensitivity and specificity are low, however, so it is of more use for monitoring than for screening or diagnosis. CEA levels are useful in assessing prognosis (with other factors), detecting recurrence (especially for disease that cannot be evaluated by other means) and monitoring treatment in people with colorectal cancer. CEA is particularly recommended for postoperative follow-up of patients with stage II and III colorectal cancer if further surgery or chemotherapy is an option, Sturgeon *et al.*, (2009).

Carcinoembryonic antigen (CEA) is one of the most widely used tumor markers worldwide. Its main application is mostly in gastrointestinal cancers, especially in colorectal malignancy.

Serial CEA measurements can detect recurrent colorectal cancer with a sensitivity of 80%, a specificity of 70%, and can provide a lead time of 5 months. CEA is the most frequent indicator of recurrence in asymptomatic patients and currently is the most cost-effective test for the preclinical detection of resectable disease. Overall, however, little evidence is available that

monitoring of all patients with diagnosed colorectal cancer leads to enhanced patient outcome or quality of life.

2.5.2 Normal range of CEA level

Individual laboratory normal ranges vary but CEA level is usually deemed to be normal at 2.5-5 μg/L. increasing levels of CEA suggestive of active disease may be more clinically helpful than absolute level, Sturgeon *et al.,* (2008). Levels exceeding 10 μg/L are rarely due to benign disease or the moderate elevation that may occur due to smoking, Fakih *et al.,* (2006).

2.5.3 Sensitivity of CEA

Sensitivity in early-stage colorectal cancer is very low and increases with the stage of the disease. Studies reporting sensitivity vary in the cut off for what constitutes a normal CEA level so results are varied. Using a cutoff point of 5μg/L, the proportions of patients with increased values are 3%, 25%, 45% and 65% for patients with Dukes' A, B, C and D disease respectively. Around 72% of cases with unresectable or metastatic disease have elevated CEA levels Sturgeon *et al.,* (2008).

2.5.4 Monitoring treatment:

The major role for CEA levels is in following patients for relapse after intended curative treatment of colorectal cancer. CEA levels typically return to normal within four to six weeks after successful surgical resection. The CEA level can also be used to assess the response to chemotherapy. It is recommended that people with stage II or III colorectal cancer have CEA surveillance every three months for at least three years after treatment. National Institute for Health and Care Excellence (NICE) guidelines recommend at least every six months, although most other authorities recommend every three months, Sturgeon *et al.,* (2008). Clinical trials

have shown improvement in survival after five years in patients who underwent CEA monitoring as part of post-treatment management, Sturgeon *et al.,* (2008); Jeffery *et al.,* (2007). However, normal levels do not necessarily indicate that recurrence has not occurred, Hara *et al.,* (2008).

Currently, the most useful application of CEA is in the detection of liver metastasis from colorectal cancers. Because of the relative success of surgery in resecting hepatic metastases, serial determinations of the marker are recommended for detecting cancer spread to the liver. There are plans that in the future, preoperative concentrations of CEA may be included with the standard staging procedures for assessing prognosis.

According to the 2001 American Association for Clinical Chemistry, Carcinoembryonic antigen (CEA) was first described in 1965 by Gold and Freedman, when they identified an antigen that was present in both fetal colon and colon adenocarcinoma but that appeared to be absent from healthy adult colon. Because the protein was detected in only cancer and embryonic tissue, it was given the name carcinoembryonic antigen, or CEA. Subsequent work showed that CEA, or at least a CEA-like molecule, was also present in certain healthy tissues, although concentrations in tumors were on average 60-fold higher than in the nonmalignant tissues, Boucher *et al.,* (1989). In one of the first reports on CEA in serum. Thomson *et al.,* (1991) found increased concentrations in 35 of 36 patients with colorectal cancer. In contrast, high values were not found in “normal” subjects, pregnant women, patients with no gastrointestinal cancers, or in patients with miscellaneous benign gastrointestinal diseases. Although these findings were not confirmed, they nevertheless prompted widespread use of carcinoembryonic antigen as a marker for colorectal cancer. Thirty years after its initial detection in serum, carcinoembryonic antigen is

one of the most widely used tumor markers worldwide and certainly the most frequently used marker in colorectal cancer.

2.5.5 Carcinoembryonic Antigen as a Marker for Colorectal Cancer screening

In screening for colorectal cancer, the aim should be to detect disease at either Dukes' A or B stage. Malignancy detected at more advanced stages is unlikely to be more treatable than that detected through the usual course of events. Using an upper limit of normal of 2.5 mg/L, (Fletcher *et al.*,1986) calculated that carcinoembryonic antigen has a sensitivity of 36% and a specificity of 87% in screening for Dukes' A and B colorectal cancer. These findings, combined with the low heterogeneity, which appears to be attributable to variations in its carbohydrate side chains, Thomas *et al.,* (1990). Most of the carbohydrate is composed of mannose, galactose, *N*acetylglucosamine, fucose, and sialic acid, Thomas *et al.,* (1990).

Carcinoembryonic Antigen is a member of the immunoglobulin superfamily. Two types of immunoglobulin domains are found: an N-terminal domain of 108 amino acids homologous to the immunoglobulin variable domain (IgV-like) and six domains homologous to the immunoglobulin constant domain of the C-2 set (IgC2-like), Thompson *et al.*, (1991; Hammerstron *et al.,* (1999).

Carcinoembryonic antigen is attached to the cell membrane by a glycosyl phosphatidylinositol anchor and probably is released as a soluble form by a phospholipase C or phospholipase D, Thompson *et al.,* (1991). Structural similarity of CEA to certain immunoglobulin-related

proteins, such as ICAM-1 and ICAM-2, initially suggested that CEA might act as an adhesion molecule. In vitro experiments showed that CEA was capable of both homophilic (CEA binding to CEA) and heterophilic (CEA binding to non-CEA molecules) interactions. Because alterations in cell adhesion are causally involved in cancer invasion and metastasis, it was further suggested that CEA may play a role in these processes, Jessup *et al.,* (1989). Evidence for a role in cancer dissemination was obtained recently by Hostetter *et al.,* (1990), who showed that after transplantation of colorectal tumors into nude mice, the number of liver metastases increased from 2% to 48% following injection of mice with CEA. There is, however, no direct evidence that CEA is causally involved in cancer dissemination.

Although *in vitro* data implicate CEA in cell adhesions, its localization to the apical surface of mature enterocytes in healthy human colon is difficult to reconcile with this role. In the healthy colon, CEA has been found to bind certain strains of *Escherichia coli*. According to Thompson *et al.*, (1991), this binding may facilitate bacterial colonization of the intestine. On the other hand, suggested that CEA may play a role in protecting the colon from microbial infection, possibly by binding and trapping infectious microorganisms.

2.6 Immunohistochemistry

Immunohistochemistry (IHC), or immunocytochemistry, is a method for localizing specific antigens in tissues or cells based on antigen-antibody recognition; it seeks to exploit the specificity provided by the binding of an antibody with its antigen at a light microscopic level. IHC has a long history, extending more than half a century from 1940, when Coons developed an immunofluorescence technique to detect corresponding antigens in frozen tissue sections. However, only since the early 1990s has the method found general application in surgical

pathology, Gown (2002). A series of technical developments led eventually to the wide range of IHC applications in use today. The enzymatic label (horseradish peroxidase), developed by Avrameas and by Nakane and colleagues (1966), allowed visualization of the labeled antibody by light microscopy in the presence of a suitable colorogenic substrate system. In Oxford, Taylor and Burns developed the first successful demonstration of antigens in routinely processed formalin-fixed paraffin-embedded (FFPE) tissues, Coppola *et al.,* (2011*).* A critical issue in the early development of immunoperoxidase techniques was related to the need to achieve greater sensitivity. Greater sensitivity would facilitate staining of FFPE tissues—from a simple one-step direct conjugate method to multiple-step detection techniques such as the peroxidase antiperoxidase (PAP), avidin-biotin conjugate (ABC), and biotinstreptavidin (B-SA) methods—and would eventually lead to amplification methods (such as tyramide) and highly sensitive " polymer-based" labeling systems, Ager *et al.*, (2015).

2.6.1 Basic Principles of Immunohistochemistry

Surgical pathologists have long recognized their fallibility, although they have not always publicized it, Gown (2002). They have, however, sought more certain means of validating morphologic judgments. A variety of "special stains" were developed to facilitate cell recognition and diagnosis; most of these early stains were based on chemical reactions of cell and tissue components in frozen sections (histochemistry). In certain circumstances, these histochemical stains proved to be of critical value in specific cell identification. More often, they served merely to highlight or emphasize cellular or histologic features that supported a particular interpretation without providing truly specific confirmation. When the new field of immunohistochemistry was created by combining immunology with histochemistry, a wide variety of truly specific special stains were generated. The aims of IHC are akin to those of

histochemistry. Indeed, IHC builds on the foundations of histochemistry; it does not replace histochemistry but rather serves as a valuable adjunct that greatly extends the variety of tissue components that can be demonstrated specifically within tissue sections or other cell preparations. As emphasized by pioneers in this field of functional morphology, "the object of all staining is to recognize microchemically the existence and distribution of substances which we have been made aware of macrochemically, Nakane *et al*, (1966).

The basic critical principle of IHC, as with any other special staining method, is a sharp visual localization of target components in the cell and tissue, based on a satisfactory signal-to-noise ratio. Amplifying the signal while reducing non-specific background staining (noise) has been a major strategy to achieve a satisfactory result that is useful in daily practice. After more than two decades, advances in IHC have provided a feasible approach to performing immunostaining on routinely processed tissues, such that this method is now "routine" for the performance of IHC "special stains" in surgical pathology laboratories using FFPE tissues. However, demands for improved reproducibility and for quantification have led to a growing recognition that IHC has the potential to be more than just a special stain. If properly controlled in all aspects of its performance, IHC can provide a tissue-based immunoassay with the reproducibility and quantitative characteristics of an ELISA (enzyme-linked immunosorbent assay) test, which not only detects the presence of an "analyte" (protein or antigen) but also provides an accurate and reliable measure of its relative or real amount.

2.6.2 Principles of Immunohistochemistry

Immunohistochemistry is been widely used in diagnosing carcinogenesis and cancer because it takes advantage of the principle of the antigen-antibody affinity reaction through its ability to

identify and localize proteins of interest via detection with labeled conjugates Rekardson (2014). This antigens have corresponding antibodies produced by the body to form a complex compounds which are usually picked up by immunochemical dyes or stains giving a distinctive color (brownish color in the case of the MLH1 gene) known as "expression.

2.7 Animal Models in Colon Carcinogenesis

2.7.1 Rodent Colon Tumor Models (History and Overview)

The study of experimental colon carcinogenesis in rodents has had a remarkably long history, dating back almost 80 years. Perhaps the earliest published study of Lorenz *et al.* (1941) demonstrated tumorigenesis in the forestomach and intestine of mice following feeding with the polycyclic aromatic hydrocarbon, methylcholanthrene. Lisco *et al.* (1947) reported that feeding radioactive yttrium to rats induced a high proportion of colon tumors. Walpole *et al.* (1952) reported that white rats given injections of 4-aminodiphenyl and 3,2-dimethyl-4-aminodiphenyl developed colon tumors. However, the most commonly used model for sporadic colorectal cancer (CRC) takes advantage of the organotropism of the colon carcinogens, 1,2-dimethylhydrazine (DMH) and azoxymethane (AOM). Identification of these carcinogens arose initially from an early population study of Guamanians in which hydrazines were reported to be possible colon carcinogens with the hydrazine source being consumed cycad flour, Laqueur *et al.*, (1963). When large quantities of cycad flour were fed to rats, adenocarcinomas in the colon arose in some of the animals. The carcinogen in cycad flour was subsequently found to be cycasin, a form of methylazoxymethanol (MAM), Laqueur *et al*., (1964). DMH, a metabolic precursor of MAM, was used in several early studies to induce tumors in rats, Thumherr *et al.*, (1975). Repetitive treatment with this methylating agent was reported to produce colon tumors in

rodents that exhibit many of the pathological features associated with the human disease, Ward, (1974). Thus, DMH has provided cancer researchers with a reproducible experimental system for studying 'sporadic' (nonfamilial) forms of CRC, LaMont, (1978*)*. However, AOM offers advantages over DMH, including enhanced potency and greater stability in dosing solution, Neufert *et al*., (2007). Since then, several thousand studies using AOM have been published Toth *et al.,* (1976). We have used this mouse model extensively for 15 years and have compiled considerable data on sensitivity and lesion characteristics across a number of mouse lines, Flynn and Kralj-Hans *et al*., (2007).

The occurrence of spontaneously occurring cancers within the gastrointestinal tract of rodents is not as rare as might be expected. For example, Rowlatt *et al.,* (1969) described the occurrence of a wide range of tumors of the intestinal tract in aged C57BL/6 mice, although the majority of the lesions were located in the small intestine. Miwa *et al.* (1976) reported 28 cases of spontaneous colon tumors in male wistar rats. Vandenberghe *et al.* (1985) in a study of Campylobacter-like bacteria reported 17 naturally occurring primary endophytic adenonocarcinomas in the ascending colon of Wistar rats. An examination of animal husbandry also demonstrated the presence of adenocarcinomas in the ascending colon of ACI rats foster bred by an Osaka female rat, perhaps depending on milk factor and humidity, Miyamoto *et al*., (1991). Newmark *et al.,* (2001) found that maintaining mice on a semipurified diet, referred to as the new Western diet, designed to recapitulate a human Western diet, was capable of inducing colon tumors in 42% of 18-month-old C57BL/6 mice. Richie *et al*., (2007) showed an induction of colon tumors in p53 knockout mice by causing glutathione depletion with buthionine sulfoximine treatment.

In summary, there are a number of advantages to studying the pathogenesis of carcinogen-induced colon cancer in rodent models. The models are highly reproducible, they can be readily tested on animals with different genetic backgrounds and the pathogenesis recapitulates human CRC, at least at the early stages. Historically, the majority of colon carcinogenesis studies have been carried out in rats, Madera *et al.* (1983). However, the high frequency of tumors generated within the distal colon of mice, as well as the histogenesis of multiple adenomas with subsequent development of adenocarcinomas, also validate the importance of this species for studying the pathogenesis of colon cancer, Nambiar *et al*., (2003). An advantage of murine models is the availability of extensive genetic information on individual mouse lines, the existence of recombinant inbred mouse panels and the ever-increasing number of transgenic, knockout and knock-in genetic models that are available for study.

2.8 Role of MNU in Colon Carcinogenesis Studies

N-methyl-N-nitrosourea (MNU), are among the primary causes of cancer. Verma *et al.,* (2013) MNU an alkylating agent was reported to induce retinal degeneration in mice, rats and other animals following systemic administration, providing a simple model of retinal degenerative diseases, Neufert *et al.,* (*2007).* It exhibits its toxicity by transferring its methyl group to nucleo bases in nucleic acids which can lead to AT: GC transition mutation, Swanson *et al.* (1996). The molecular mechanism of MNU involves specific G-35 point mutation in codon 12 which results in substitution of normal glycine with an aspartic acid, Saminathan *et al*., (2014). A clear example of MNU toxicity is seen when cells are exposed to the carcinogen that leading to inactivation of genes required for repair of base-base mismatches in DNA, referred to as mismatch repair (MMR) genes (MLH1, MLH2, etc). This damaged or deactivated genes then

lacks the ability to repair DNA mutation which then accumulates and prompts cells to replicate uncontrollably, This MMR deficiency can be detected during Immunohistochemical analysis. Exposure to MNU can result in skin and eye irritation headache, nausea, and vomiting. MNU is reasonably anticipated to be a carcinogen based on sufficient evidence of carcinogenicity in experimental animals.

Various cancers which were successfully induced in animal models include: squamous cell carcinomas of the fore-stomach, sarcomas and gliomas of the brain, adenocarcinomas of the pancreas, mammary carcinomas, leukemia, and lymphomas, Perse *et al*., (2009).

2.9 Haematological Parameters in Cancer/Colon Carcinogenesis and their Application as Possible Cancer Biomarker.

Haematology refers to the study of the numbers and morphology of the cellular elements of the blood, the red cells (erythrocytes), white cells (leucocytes), and the platelets (thrombocytes) and the use of these results in the diagnosis and monitoring of disease, Merck Manual, (2012). Haemtological studies are useful in the diagnosis of many diseases as well as investigation of the extent of damage to blood, Onyeyili *et al.* (1992)*;* Togun *et al.*, (2007). Haematological studies are of ecological and physiological interest in helping to understand the relationship of blood characteristics to the environment, Ovuru and Ekweozor (2004) and so could be useful in the selection of animals that are genetically resistant to certain diseases and environmental conditions, Mmereole (2008); Isaac *et al.* (2013). Haematological parameters are good indicators of the physiological status of animals, Khan and Zafar, (2005). Haematological parameters are those parameters that are related to the blood and blood forming organs, Waugh *et al.* (2001)*;* Bamishaiye *et al*. (2009). Blood act as a pathological reflector of the status of exposed animals

to toxicant and other conditions, Olafedehan *et al.,* (2010). As reported by Isaac *et al.* (2013), animals with good blood composition are likely to show good performance.

Laboratory tests on the blood are vital tools that help detect any deviation from normal in the animal or human body, Ogunbajo *et al.* (2009). The examination of blood gives the opportunity to investigate the presence of several metabolites and other constituents in the body of animals and it plays a vital role in the physiological, nutrition and pathological status of an organism, Aderemi (2004); Doyle (2006). According to Olafedehan *et al.* (2010), examining blood for their constituents can provide important information for the diagnosis and prognosis of diseases in animals. Blood constituents change in relation to the physiological conditions of health (Togun *et al.,* 2007). These changes are of value in assessing response of animals to various physiological situations, Khan and Zafar (2005). According to Afolabi (2010), changes in haematological parameters are often used to determine various status of the body and to determine stresses due to environmental, nutritional and/or pathological factors.

2.9.1 Haematological Components and their Functions

Blood which is a vital special circulatory tissue is composed of cells suspended in a fluid intercellular substance (plasma) with the major function of maintaining homeostasis Soetan *et al.,* (2013). Haematological components, which consist of red blood cells, white blood cells or leucocytes, mean corpuscular volume, mean corpuscular haemoglobin and mean corpuscular haemoglobin concentration are valuable in monitoring feed toxicity especially with feed constituents that affect the blood as well as the health status of farm animals, Oyawoye and Ogunkunle (2004). Red blood cells (erythrocytes) serve as a carrier of haemoglobin. It is this haemoglobin that reacts with oxygen carried in the blood to form oxyhaemoglobin during

respiration Chineke *et al.,* (2006). According to Soetan *et al.,* (2013) red blood cell is involved in the transport of oxygen and carbon dioxide in the body. Thus, a reduced red blood cell count implies a reduction in the level of oxygen that would be carried to the tissues as well as the level of carbon dioxide returned to the lungs, Ugwuene (2011); Soetan *et al.,* (2013).

The major functions of the white blood cell and its differentials are to fight infections, defend the body by phagocytocis against invasion by foreign organisms and to produce or at least transport and distribute antibodies in immune response. Thus, animals with low white blood cells are exposed to high risk of disease infection, while those with high counts are capable of generating antibodies in the process of phagocytocis and have high degree of resistance to diseases, Soetan *et al.,* (2013) and enhance adaptability to local environmental and disease prevalent conditions, Kabir *et al.,* (2011); Okunlola *et al.* (2012).

Blood platelets are implicated in blood clotting. Low platelet concentration suggests that the process of clot-formation (blood clotting) will be prolonged resulting in excessive loss of blood in the case of injury. Packed Cell Volume (PCV) which is also known as haematocrit (Ht or Hct) or erythrocyte volume fraction (EVF), is the percentage (%) of red blood cells in blood (Purves *et al.* 2003). According to Soetan *et al.* (2013) packed cell volume is involved in the transport of oxygen and absorbed nutrients. Increased packed cell volume shows a better transportation and thus results in an increased primary and secondary polycythemia. Haemoglobin is the iron-containing oxygen-transport metalloprotein in the red blood cells of all vertebrates, Maton *et al.,* (1993) with the exception of the fish family, channichthyldae, Sidell and Brien, (2006) as well as tissues of invertebrates. Haemoglobin has the physiological function of transporting oxygen to

tissues of the animal for oxidation of ingested food so as to release energy for the other body functions as well as transport carbon dioxide out of the body of animals, Ugwuene (2011).

2.10 Oxidative Stress and Cancer / Colon Carcinogenesis.

Oxidative stress, defined as the imbalance between oxidants and antioxidants in favor of the oxidants potentially leading to damage has been suggested to be that which cause aging and various diseases in humans. In modern western medicine, the balance between anti-oxidation and oxidation is believed to be a critical concept of maintaining a healthy biological system. Oxidative stress is among the major causative factors in the induction of many chronic and degenerative diseases including atherosclerosis, diabetes mellitus, cancer, Parkinson's disease, immune dysfunction and ageing, Souri *et al*., (2008). Antioxidants can be effective in the prevention of free radical formation by scavenging or promotion of their decomposition and thus, suppress such disorders. Hence, there is growing interest in natural antioxidants from herbal and food plant sources, Atawodi (2010); Ghanbari *et al.*, (2012) as epidemiological and laboratory studies have strongly supported their therapeutic and protective efficacies in biological systems, Souri *et al.,* (2008); Atawodi (2011). Besides, vegetables, spices and herbs are rich in natural antioxidants that protects against oxidative stress and thus play important roles in the therapy and chemoprevention of diseases that have their etiology and pathophysiology in reactive oxygen species, Atawodi *et al.,* (2010); Manzoor *et al.,* (2012).

Oxidative DNA damage is a major source of the mutation load in living organisms, with more than one hundred oxidative DNA adducts (purine, pyrimidine, and the deoxyribose backbone) having been identified, Lu *et al.,* (2001); von Sonntag (1987); Dizdaroglu (1992); Demple and Harrison (1994). The estimated frequency of oxidative DNA damage in human cells is 104 lesions/cell/day, Fraga et al. (1990); Lu *et al.,* (2001). Being highly reactive, the hydroxyl radical

is the predominant ROS that targets DNA, Lu *et al.,* (2001). Hydrogen peroxide, a precursor to hydroxyl radical, is less reactive and more readily diffusible and thus more likely to be involved in the formation of oxidized bases through Fenton and Haber-Weiss reactions, Guyton and Kensler (1993); Barber and Harris (1994). ROS induced DNA damage can result in single- or double-strand breakage, base modifications, deoxyribose modification, and DNA cross-linking. Cell death, DNA mutation replication errors, and genomic instability can occur if the oxidative DNA damage is not repaired prior to DNA replication, Marnett (2000); J. P. Cooke (2003); Klaunig and Kamendulis (2004); Valko *et al.,* (2006).

In traditional Chinese medicine for more than 2000 years, a general recommendation to their consumption is to increase the intake of foods rich in antioxidant compounds (e,g polyphenols, carotenoids) due to their well-known healthy effects. As a consequence, these evidences accelerated the search for antioxidants principles, which led to the identification of natural resources and isolation of active antioxidant molecules. Many plants have been identified as having potential antioxidant activities and their consumption recommended, Kitts, *et al.,* (2000); Lee *et al.,* (2003).

Bioactive phenols are very interesting as antioxidants because of their ability to act as efficient free radical Abbreviations: BHT, Butylated hydroxytoluene; DPPH, 1,1- diphenyl-2-picrilhydrazyl; AML, acute myeloid leukemia, Langley-Evans, (2000). In the last two decades the number of publications on the potential health benefits of polyphenols, have increased enormously, Tiwari, (2001); Katalinic *et al.,* (2006).

2.11 Antioxidants

Antioxidants are believed to play a very important role in the body defense system against ROS Boxin et al. (2002), Vivek and Surendra (2006). In another term antioxidant is "any substance that, when present at low concentrations compared with that of an oxidizable substrate, significantly delays or inhibits oxidation of that substrate, Halliwell and Gutteridge (1995). Halliwell (2007) reported that an antioxidant is "any substance that delays, prevents or removes oxidative damage to a target molecule. Antioxidants are an inhibitor of the process of oxidation, even at relatively small concentration and thus have diverse physiological role in the body. Antioxidant constituents of the plant material act as radical scavengers, and helps in converting the radicals to less reactive species. A variety of free radical scavenging antioxidants is found in dietary sources like fruits, vegetables and tea, etc

Natural products of plant origin are major source of vitamin C, vitamin E, and beta –carotene and have been proposed as potential source of natural antioxidants with strong activity (Block *et al.*, 1992). This activity is mainly due to the presence of phenolic compounds such as flavonoids, phenols, flavonols and proanthocyanidins, Bose *et al.*, (2012). Antioxidants are chemical compounds which contain monohydroxy/polyhydroxy phenol; they just work to slow down the lipid peroxidation, German *et al.*, (2013). These compounds have low activation energy to donate hydrogen atom and therefore, cannot initiate the second free radicals. The free radical electrons are stable and thus, slow down the oxidation. Cells contain many antioxidant systems to prevent injury. Prevention of excessive ROS and repair of cellular damage is essential for cell's life, Guttridge *et al.*, (2011).

Polyphenols are the major plant compounds with high level of antioxidant activity. This activity could be due to their ability to absorb, neutralize and to quench free radicals, Bose *et al.*, (2012). Their ability as free radical scavenger could also be attributed to their redox properties, presence

of conjugated ring structures and carboxylic group which have been reported to inhibit lipid peroxidation, Bose *et al.,* (2012). Antioxidant is one of the most essential ingredients of today's therapy since they reduce in vivo oxidative damages. Plants are the good resources for natural antioxidants. DPPH radicals scavenging method is widely used to investigate the total antioxidant activities in plants. Antioxidant substances block the action of free radicals which have been implicated in the pathogenesis of many diseases including atherosclerosis, ischemic heart disease, cancer, Alzheimer's disease and in the aging process.

In addition to dietary antioxidants, the body relies on several endogenous defense mechanisms to help protect against free radical-induced cell damage. The antioxidant enzymes – glutathione peroxidase, catalase, and superoxide dismutase (SOD) – metabolize oxidative toxic intermediates and require micronutrient cofactors such as selenium, iron, copper, zinc, and manganese for optimum catalytic activity, Anuj *et al.,* (2016).

Superoxide dismutase (SODs) are a group of key enzymes functioning as the first line of antioxidant defense by virtue of the ability to convert highly reactive superoxide radicals (dismutation) into hydrogen peroxide and molecular oxygen, Wang *et al.,* (2016).

Catalase is a tetrameric porphyrin-containing enzyme that is located mainly in peroxisomes. It catalyzes the conversion of H2O2 to water and molecular oxygen in two steps, Valko *et al.,* (2006); Chaudiere *et al.,* (1999), and Aslani *et al.,* (2016).

Catalase-Fe (III) + H2O2→compound I

Compound I + H2O2→catalase-Fe (III) + 2H2O + O2

Ligand induced ROS is generated in nonphagocytic cells, Fanburg *et al.,* (2015). NADPH oxidase is activated in vascular smooth muscle cells by Angiotensin II and 5-HT neurotransmitter. This in turn, generates hydrogen peroxides and superoxide anions. ROS activated by Ang-II stimulate the p38 Mitogen Activated Protein Kinase (MAPK) pathway and cause cell hypertrophy. ROS activated by 5-HT stimulate ERK (extra cellular signal regulated kinase), MAPK pathway and cause cell proliferation.

2.1.2 Inflammation

Inflammation is a protective strategy evolved in higher organisms in response to detrimental insults such as microbial infection, tissue injury and other noxious conditions. It is an essential immune response by the host that enables the removal of harmful stimuli as well as the healing of damaged tissue. Acute inflammation has therefore been considered as a part of innate immunity, the first line of host defense against foreign invaders and danger molecules. Mankind has known the classical symptoms of inflammation for hundreds of years, which include redness, pain, swelling and heat, Medzhito**v** (2008).

As a protective strategy for the host, one of the main aims of inflammation is to reinstate cellular homeostasis in response to any damaging condition. The mechanism underlying the initiation of inflammation is therefore tightly connected to the physiological state of homeostasis. And so, inflammation is considered as an 'adaptive response' to any harmful effect threatening the integrity of the cellular homeostasis. It is quite plausible to understand that such an adaptive response operates at the expense of normal cellular functions, Medzhitov (2010). As a result, the

longer this response persists in, the host will encounter the more damaging consequences. In contrary to its beneficial role as a safeguard for cellular physiology, the inflammatory response is required to be least enduring in order to avoid any escalation of its unfavorable circumstances.

2.1.2.1 Molecular mechanism underlying inflammation

Inflammatory stimuli are first recognized by the host cells through specific transmembrane receptors, called pattern recognition receptors (PRRs), which are expressed by cells of both innate and adaptive immune systems. PRRs are germline-encoded receptors, which are responsible for sensing the presence of infecting microorganisms as well as the incidence of any cellular damage (Akira et al., 2006). They do so by recognizing structures conserved in microbes, called pathogen-associated molecular patterns (PAMPs), as well as endogenous molecules derived from internal injuries, called danger-associated molecular patterns (DAMPs). To date, a number of PRRs have been identified with the selective ability to detect PAMPs, DAMPs or both and these include Toll-like receptors (TLRs), C-type lectin receptors (CLRs), RIG-1-like receptors (RLRs) and NOD-like receptors (NLRs). The interactions of these receptors with the appropriate stimuli result in transmitting signals to nucleus where the activation of a selective set of genes takes place via both transcriptional and posttranscriptional mechanisms, Akira *et al.*, (2006); Medzhitov (2007).

2.1.2.2 Inflammation and cancer

Over the last decade it has become evident that inflammation plays a critical role in promoting cancer, in particular the tumorigenesis, a process of tumor formation. In addition to cancer cells, various types of immune cells are commonly found within tumors. Interestingly, an inflammatory microenvironment is also more frequently found as an essential part of all tumors,

Mantovanietal (2008); de Visseretal (2006). It has been demonstrated that the inflammatory response triggered by infection is also associated with increased cancer risk, de Martel and Franceschi (2009). A recent study has revealed that lung tumorigenesis caused by tobacco smoke is in fact initiated by IKKbeta and JNK1-mediated chronic inflammation, suggesting that the pathways of both tumorigenesis and inflammation are closely linked, Takahashietal (2010). It is believed that apart from the tumor-promoting inflammatory response, an active anti-tumor immunity is also present in most tumor microenvironments. It is thus suggested that the progression of tumorigenesis is dictated by the battle between tumor-promoting inflammation and anti-tumor immunity. Obviously, in established tumors, antitumor immunity is profoundly dominated by tumor-promoting inflammation, Smyth *et al.,* (2006); Lin and Karin (2007). Studies show that inflammatory microenvironments not only promote cancer cell growth but also increase mutation rates, possibly by producing reactive oxygen species and nitrogen intermediates which can cause DNA damage and genomic instability, Grivennikov *et al.,* (2010). The role of inflammation in genomic instability is supported by the observation that activation-induced cytidine deaminase (AID), an enzyme which triggers genomic instability and is usually overexpressed in many cancers, is induced by inflammatory cytokines Okazaki *et al.,* (2007). Apart from genomic instability, environmental factors such as carcinogens, infectious microbes, tobacco smoke and inhaled pollutants have been considered to play critical roles in inflammation induced cancers, Aggarwal *et al.,* (2009). Therefore, not all chronic inflammatory diseases are connected to cancer risk. For example, rheumatoid arthritis does not promote cancer whereas inflammatory bowel disease and chronic hepatitis do due to the exposure to dietary and environmental carcinogens. The connection between inflammation and cancer does not operate in one direction only as numerous studies showed that DNA damage can also lead to

inflammation. Using the model of hepatocellular carcinoma induced by the carcinogen diethyl nitrosamine (DEN), several reports demonstrated that the DNA damage-induced necrotic cell death lead to inflammation, Maeda *et al.,* (2005); Sakurai *et al.,* (2008). Cancer-associated oncoproteins such as Ras, Myc and RET can also lead to inflammation by activating signaling pathways involved in the production of pro inflammatory cytokines and chemokines, Mantovani *et al.,* (2008). Last of all, tumor-associated inflammation can also be instigated by the modern cancer therapy itself such as radiotherapy and chemotherapy. These therapies are usually associated with a substantial amount of necrotic death of not only cancer cells but also surrounding normal cells, which in turn activates the inflammatory response, Zong and Thompson (2006). Therefore, based on the literature, it can be concluded that inflammatory response is an integrated part of cancer biology, right from the beginning of tumorigenesis toward the application of therapeutic interventions.

2.12 *Cucurbita Species*

2.12.1 Description of plant:

Taxonomical Position of *Cucurbita maximus*

Division:	Spermatophyta
Sub-Division:	Angiospermae
Class:	Dicotyledonae
Sub-Class:	Polypetalae
Series:	Caliciflorae
Order:	Passiflorales
Family:	Cucurbitaceae
Genus:	*Cucurbita*
Species:	*maximus* **Source**: International Journal of Phytopharmacy

Pumpkin leaves are orbicular in outline and not pointed or much lobed, with deep sinus at base. In general, pumpkin stems are more rigid, prickly, and angular (with an approximate five-degree angle) than squash stems, which are generally softer, more rounded, and more flared where joined to the fruit.

The pumpkin varies greatly in shape, ranging from oblate to oblong. The rind is smooth and usually lightly ribbed. Although pumpkins are usually orange or yellow, some fruits are dark green, pale green, orange-yellow, white, red and gray. Leaves are nearly orbicular in outline and not pointed or much lobed but with deep sinus at base. Corolla prevailingly with soft abtuse more or less crinkly revolute or hanging lobes, the tube with parallel sides or bulging at base, peduncle short, spongy, nearly cylindrical, not expanded at attachment to fruit, (Lirtika *et al.*,2011).



Plate 2.1: Picture of the fruit sizes, shapes and color for *Cucurbita maxima* which were collected from Ochi Ogbonicha in Ofu LGA Kogi State, Nigeria.

Cucurbita maxima is a guard like squash belonging to the genus *cucurbita* and the family curcubitaceae. Other species in this genus include *Curcubita mixta*, *Curcubita pepo* and

Curcubita moschota Singh (2012). It is known by various names: for instance, it is known as Giant pumpkin or winter squash in English, Echi in Igala Language, Kabewa in Hausa, ebeshe in Nupe, Ogbokolo, Okoro, Anya in Igbo and Elegede, Isi, in Yoruba. It is collectively ranked among the 10 leading vegetable crops worldwide. China and India are the world leading producers. Other major producers are U.S.A, Egypt, Mexico, Ukraine, Cuba, Italy, Iran and Turkey (FAOSTAT, 2007). Squashes (*Cucurbita* spp.) are members of the economically important *Cucurbitaceae*. There are three economically important *Cucurbita* species, namely *Cucurbita pepo* L., *Cucurbita maxima* Duch and *Cucurbita moschata* Duch, which have different climatic adaptations and are widely distributed in agricultural regions worldwide, Paris and Brown (2011); Wu *et al.,* (2007).

2.13 Origin and Distribution

The *Cucurbita* genus is of American origin. *Cucurbita maxima* are the most diverse *Cucurbita* species after *Cucurbita pepo* and are found throughout tropical and temperate regions. Phenotypic diversity within populations of *Cucurbita* is high and includes variation in shape, size and colour of fruits; number and size of seeds; quality, color and thickness of fruit flesh and precocity in fruit production, among other traits, Whitaker and Robinson, (1986); Hernandez *et al*., (2012). The centre of diversity, *Cucurbita maxima*, lies in South American temperate zones, where landraces exhibit an array of interesting traits. Many landraces of this species are also found in North America, Australia and different countries of Africa **(**Zambia and Nigeria**)**; Asia (China, India, and Iran) and Europe (Spain and Turkey), Ferriol and Pico (2008). Turkey is one of the important diversity centers for cultivated cucurbits because of their adaptation to diverse ecological conditions as a result of natural selection and also farmers' selections, Sari *et al*., (2008). Pumpkins are native to North and Central America, Whitaker (1975), where some

Cucurbita varieties were cultivated before Europeans reached that area. Nowadays, pumpkins are cultivated in almost all areas with an appropriate climate in the world.

2.14 Culinary Uses

The fruits are the most valuable part of *Cucurbita* plants; some of them own a high nutritional value and a high level of carotenoids and for this reason, in many countries, *Cucurbita* fruits are important sources of provitamin A in human nutrition, being consumed usually boiled or baked, according to the local traditions, Muntean *et al.,* (2013). Almost all parts of *Cucurbita maxima* plants are edible and healthful. The fruits (pumpkins or squashes) are well known; they are large, weighing 8 to 10 kg, sometimes up to 20 kg. The orange and yellow color of their flesh is related with the contained carotenoids – phytochemicals that may reduce risk of some cancers, heart attacks and some aging processes, Muntean *et al.*, (2013). The flowers are also loaded with carotenoids and can be eaten raw in salads or cooked. The leaves contain lot of iron and other minerals and vitamins; smaller new leaves are good ingredients in salads, while stems can be chopped and boiled with other vegetables, providing fibers. The seeds can be dried and eaten raw or baked; they contain valuable nutrients oils, proteins and vitamins, Muntean *et al.*, (2013).

2.15 Medicinal Uses

Pumpkin is a creeper and is extensively found very frequently on the roofs of houses all over India. The parts which are used are as follows, seed, pulp, and fruit stalk. The fruits of pumpkin are large weighing 8 to 10 kg, sometimes up to 20kg.the medium sized fruit of "Arka Chandan", an old variety from Indian institute of horticulture research, Hessaraghatta, weighs 2 to 3 kg and has orange color flesh.

Pumpkin seeds contain many valuable functional components and have been traditionally used for herbal, therapeutic as well as clinical application. Pumpkin seeds have been used as safe deworming and diuretic agents, and the seed oil as a nervine tonic, Das *et al.,* (2010). Pumpkin seed oil has a strong antioxidant property, and has been recognized for several health benefits such as prevention of the growth and reduction of the size of prostate, reduction of bladder and urethral pressure and improving bladder compliance, alleviation of diabetes by promoting hypoglycemic activity, and lowering level of gastric, breast, lung, and colorectal cancer. Pumpkin is cultivated throughout India and in most warm region of the world, Attarde *et al*., (2010).

2.16 Chemical Composition

The chemical constituent such as 24 β ethyl 5 α cholesta-7, 22, 25- trien-3 β ol, 24 β-ethyl-5α-cholesta-7, 25-dien-3 β ol, avenasterol, spinasterol, 24 dihydrospinasterol, 24-methyl lathosterol and 25-dehydrofungisterol has been obtained from seeds, Das *et al.,* (2010). 25-dehydroporiferasterol, clerosterol, isofucosterol, stigmasterol, sitosterol, campesterol, and codisterol, isolated from seeds. 13 C-NMR of chondrillasterol studied, isolation of two new carotenoidscucubitaxanthins A and B and their structure elucidation.

Similar to other cucurbitaceous plant it contains saponin, fixed oil, resin, protein, sugar and starch. The fresh vegetable contains 89.50 % moisture and the completely dried material contains ethereal extract of albuminoids 6.12% (containing nitrogen 0.98%), soluble carbohydrates 77.33%, woody fifer 8.55%, and ash 7.00%. The seeds collected from Pakistan yields an oil (39.1%) having the following physic chemical constituents, the 3 fatty acid composition of oil is

as follows, myristic, palmitic, stearic, oleic and linoleic acid 30.3%. seeds contains sterol glycosides and sterols fatty acid esters which showed antitumor activity in mouse.

The alcoholic and etheral extracts of pumpkin seeds were found to be active against the nematode Fasciolopis buski, in vitro and in vivo test systems, Kirtika *et al.,* (2009). The therapeutic efficacy of the seeds in the treatment of nematodiasis in calves below 6 month of age was found to be 40.6% by the oral route. Chemical composition of the leaves is as follows, moisture 85.9%, lipid 1.75%, protein 4.58% and ash 1.82%.

Composition of lipid portion is as follows, nonpolar lipids 45.1%, glycolipids 24.4%, and phospholipids 30.5%, Kirtika *et al.,* (2009). The major fatty acid present in the lipid was linolenic. According to Kirtika *et al.,* (2009). The leaves contain (mg/100g) calcium 36.38%, magnesium 38.80, iron 2.04%, zinc 0.76%, copper 0.42%. An esterase was purified from the fruits.

2.17 Known Pharmacological Activities

2.17.1 Anticancer Activity- Anticancer activity of methanol extract of *Cucurbita maxima* against Ehrlich as-cites carcinoma.

Cancer is a pathological state involving uncontrolled proliferation of tumor cells. The study was carried out to investigate the antitumor potential of MECM (methanol extract of *Cucurbita maximus*) against EAC (Ehrlich Ascites Carcinoma) bearing mice. EAC is a very rapidly growing carcinoma with very aggressive behavior. It is able to grow in almost all strains of mice. The Ehrlich Ascitic tumor implantation induces *per se* a local inflammatory reaction, with increasing vascular permeability, which results in an intense edema formation, cellular

migration, and a progressive ascitic fluid formation and accumulation. The ascitic fluid is essential for tumor growth, since it constitutes a direct nutritional source for tumor cells. MECM treatment significantly reduced tumor volume probably by lowering the ascitic nutritional fluid volume. Further, the packed cell volume and the number of viable EAC tumor cells in peritoneum were significantly lower in the mice treated with MECM when compared to the tumor control group. These results could indicate either a direct cytotoxic effect of MECM on tumor cells or an indirect local effect, which may involve macrophage activation and vascular permeability inhibition, Saha, *et al.,* (2011a).

2.17.2 Antiobesity Activity - Potentiative Activity Of Cucurbita Maxima Seed Extract With Beta Vulgaris & Smilax Regelii Root Extract To Reduce Extra Fats From The Body.

The seed extract of *Cucurbita maxima* is effective for reducing fats individually as well as its combination effort through different mechanism. *Cucurbita maxima* seed sample is rich of manganese having tendency to produce thyroxine, Das *et al.,* (2010).

2.17.3 Antidiabetic Activity – Antidiabetic activity of *Cucurbita maxima* aerial parts

Streptozocin, a glucose analogue is a potent diabetogenic agent and widely used for inducing diabetes in a variety of animals by the selective degeneration and necrosis of pancreatic cells, Saha *et al.,* (2011a). The study on the Antidiabetic activity of *Cucurbita maxiam* indicates that MECM significantly reduced the elevated fasting blood glucose level with to those of diabetic control animals, Saha, *et al*., (2011b).

2.17.4 Evaluation of Hepatoprotective Activity of *Cucurbita maxima* Aerial Parts.

The study on hepatoprotective activity of Cucurbita maxima was performed to investigate the hepatoprotective activity of methanol extract of *Cucurbita maxima Duchesne* (*Cucurbitaceae*) (MECM) against carbon-tetrachloride (CCl4) induced hepatotoxicity. Liver damage induced by CCl4 is a commonly widely used model for the screening of hepatoprotective drugs. CCl4 is biotransformed by the cytochrome P- 450 system to produce the trichloromethyl free radical (CCl3‡), and this further reacts very rapidly with oxygen to yield a highly reactive trichloromethyl peroxy radical (CCl3OO‡). This free radical in turn covalently binds to cell membranes and organelle to elicit lipid peroxidation, also disturbs Ca2+ homeostasis, and finally results in cell death. Estimating the activities of serum marker enzymes, like SGOT, SGPT and ALP can make assessment of liver function. When liver cell plasma membrane is damaged, a variety of enzymes normally located in the cytosol, are released in to the blood stream. Their estimation in the serum is a useful quantitative marker of the extent and type of hepatocellular damage, Saha *et al.,* (2011e).

2.17.5 Diuretic Activity - Diuretic activity of seeds of *Cucurbita maxima* duchesne in wistar rats.

The seeds of *Cucurbita maxima* Duchesne are used traditionally as diuretics reported by Jose *et al.,* 2008. The concentration of Na+ and K+ in urine was determined by flame photometer. The volume of urine and Na+ and K+ concentration of test group was compared with the control group. The results revealed that the aqueous extract of seeds of *Cucurbita maxima* showed significant increase in urine volume when compared to control group. But the excretion of Na+ and K+ in urine was not significantly increased in drug treated group when compared to control group, Jose *et al*., (2008).

CHAPTER THREE

3.0 MATERIALS AND METHODS

3.1 Materials

3.1.1 Chemicals

The carcinogen, N-Methyl-N-Nitrosourea (MNU) was procured from BDH Chemical Ltd, Poole, England. Carcinoembryonic antigen assay kit was procured from Diagnostic Automation Inc. CA. Other chemicals used are formal-saline, graded alcohol, xylene, paraffin, eosin, hematoxylin, were obtained from the department of Histopathology, Ahmadu Bello University Teaching Hospital (ABUTH), Shika. All other reagents and chemicals were of analytical grade.

3.1.2 Feeds Used

Grower mash feed produced by UAC livestock's company was used in combination with the Powdered *Cucurbita maxima* fruit.

3.1.3 Animals used

A total of 42 Wister rats were used for the experiment

3.2 Apparatus and Equipment

Based on the experimental design, the equipment used are listed below:

Microplate reader (GM-3000, model 200711012; B. Bran Scientific Company England), pH meter (Jenway 3150 model, UK.),UV spectrophotometer (Jenway 6305 model, 50/60Hz, U.K.),

face mask, cooler, blender, pH meter (Jenway 3150 model, UK.), dissecting set, centrifuge, homogenizer, syringe, cannula, microscope, hand glove, electrical weighing balance (GF-2000, A&D Instrument Ltd, Japan), water trough, cages, feeders, microtome machine, micropipette, dissecting tape, masking tape, pins, ice blocks, tissue papers, cotton wools, nylon bag, permanent marker, dissecting set.

3.2 Methods

3.3 Sample Collection and Preparation

Fresh fruits of *Cucurbita maxima* weighing between (1kg to 2.5kg) were collected from Ochi Ogbonicha village in Ofu LGA of Kogi State Nigeria and taken to the Herbarium of the Department of Biological Sciences, Ahmadu Bello University, Zaria, for identification, authentication and a voucher specimen was deposited with number 1077. The fruits were washed in clean water, cut to remove seeds and sliced into cubes of approximately 2 mm, and later dried under morning sun (8 am to 10 am) for approximately 7 days and pounded to fine texture. The powder was kept at room temperature in an air tight container till used, and away from sunlight.

3.4 Preparation of Experimental Diets

Grower mash feed produced by UAC Livestock Company was procured from Samaru major market, Zaria Kaduna State and was used in the overall experiment. The feed served as normal diet. The powdered *Cucurbita maxima* fruit was added to the normal feed and mixed evenly to produce dietary inclusion of 0%, 2.5%, 5% and 10%. The procedure for mixing the normal feed and powdered Cucurbita maxima fruit is done by addition of percentage of fruit powder to the normal feed with addition of little distilled water (by sprinkling) to form a complete dough.

3.5 Feeding and Acclimatization of Experimental Animals

A total of forty two (42) healthy male Wistar rats weighing 80-100g were purchased from the Animal Unit of the National Veterinary Research Institute (NVRI) Vom, Jos Plateau state, Nigeria. The animals were housed in standard cages with wooden shavings as their beddings at room temperature at the Department of Biochemistry Animal House, Ahmadu Bello University Zaria and allowed free access to water and fed standard diet *ad libitum.* The animals were given two weeks of acclimatization period.

3.6. Experimental Grouping / Design

At the end of the acclimatization period, all animals were weighed and randomly divided into six (6) groups of seven animals each (n=7). The experimental animals were fed according to the group's percentage of dietary inclusion or standard diet for six (6) weeks and subsequently given MNU for period of 12 weeks and or normal saline as described below. The groups are as follows:

Group A: Normal saline + standard feed.

Group B: MNU Control: MNU + Standard feed.

Group C: MNU + dietary inclusion of 2.5% *Cucurbita maxima* powder.

Group D: MNU + dietary inclusion of 5% *Cucurbita maxima* powder.

Group E: MNU + dietary inclusion of 10% *Cucurbita maxima* powder.

Group F: Dietary inclusion control normal saline + 10% *Cucurbita maxima* powder.




Figure 3.1: Experimental Design for the Dietary Inclusion for Prevention of Colon Carcinogenesis in Male Wistar Rats.

3.6 Induction of Colon Cancer

After six (6) weeks of supplemental feeding (inclusion), the rats in the experimental groups received an intra-rectal instillation of freshly prepared MNU solution (2.0mg) using a feeding tube with a cannula while the rats in the control group also received an intra-rectal instillation of freshly prepared normal saline (0.87%. Nacl).

3.7 Collection of Blood, Organs and Tissues

The animals were sacrificed by decapitation one week after the end of the twelve weeks of the last MNU administration and the blood was collected in heparinized test tubes. The tubes for serum samples were allowed to stand at room temperature for 30 minutes, the clot was removed by centrifugation at 2000 x g for 10 minutes, the supernatant was collected using Pasteur pipette into polypropylene tubes and stored at -20^{o}c for subesequent analysis. The animals were dissected and a portion of the colon, spleen, lung, liver, and kidney were removed and fixed quickly in 10% formalin for subsequent analyses.

The colon, liver and kidney (100 mg tissue/ml buffer) were crushed using mortar and pestle and homogenized in 50mM phosphate buffer (pH 7.4); the homogenate was centrifuged at 10,000 rpm for 10 minutes and the supernatant collected with Pasteur pipette which was used for analysis. The supernatant of each organ were collected separately in a clean label plastic tube and stored at -20^{o}c until required analysis.

3.8 Lipid peroxidation by measuring the malondialdehyde (MDA) level

The concentration of thiobarbituric acid reactive substances (TBARS) in the tissue homogenate was estimated by the method of, Fraga *et al.*, (1988). The formation of malondialdehyde is the basis for the well-known TBA method used for evaluating the extent of lipid peroxidation.

Exactly 1ml of the tissue homogenate was treated with 2.0 ml of thiobarbituric acid–trichloro acetic acid–hydrochloric acid reagent and mixed thoroughly. The mixture was incubated in a boiling water bath for 15 min, cooled and centrifuged at 3000rpm for 10 min. The absorbance of the supernatant were read at 535 nm against the reagent blank.

Malondialdehyde concentration was determined using the formula below:

$$\text{Concentration (nmol/ml)} = \frac{\textbf{Absorbance of the sample}}{\textbf{E x 1.0x10}^{-9}\textbf{ x1000}}$$

Where E is molar extinction coefficient = 1.56×10^5 nM/L

3.8.1 Estimation of Stress Status / Free Radicals Generation

3.8.2 Superoxide Dismutase (SOD) activity

The enzyme superoxide dismutase (SOD) decomposes superoxide anion into hydrogen peroxide and oxygen at a high reaction rate. SOD activity assay method is based on monitoring the auto-oxidation rate of haematoxylin as originally described by Martin *et al.,* (1987). In the presence of SOD enzyme, the rate of auto oxidation is inhibited and the percentage of inhibition is linearly proportional to the amount of SOD present within a specific range. Sample SOD activity was determined by measuring ratios of auto-oxidation rates in the presence and absence of the sample.

$$\text{Hematoxylin} \xrightarrow{\text{Auto-oxidation}} H_20 + \text{Hematin (active form of hematoxylin)}$$

The procedure involved adding 920 µL of phosphate buffer (pH 7.8), which was added into clean test tube containing 40 µL of sample, mixed and incubated for 2 min at 25ºC, before 40 µL of hematoxylin was added, mixed quickly and the absorbance was measured at 560nm.

Superoxide dismutase Activity was calculated using the formula below:

$$\text{Unit of Activity (U/min/mg protein)} = \frac{\mathbf{R \times 100\ (100-R)}}{\mathbf{50 \times PC}}$$

Where $R = (Abs_B - Abs_S)$, PC= Protein concentration, Abs_B = Absorbance of Blank,

$Abs_S = Abs_B$Absorbance of Sample.

3.8.3 Catalase assay:

Catalase (CAT) activity was measured using the procedure reported by Abei (1984). Briefly, the method is as follows: 10 µL of serum was added to test tube containing 2.8 mL of 50 mM phosphate buffer (pH 7.0). The reaction was initiated by adding 0.1 ml of fresh 30 mM H_2O_2 and the decomposition rate of H_2O_2 was measured at 240 nm for 5 min. on a spectrophotometer (Jenway 640 UV/Vis). A molar extinction coefficient of 0.0411 mM^{-1} cm^{-1} was used to calculate catalase activity, using the formula below:

$$\text{Unit of Activity (U/min/mg protein)} = \frac{\mathbf{Change\ in\ Absorbance}}{\mathbf{E \times Abs_B \times PC}}$$

Where A = Absorbance of Blank, E = Molar extinction coefficient PC= Protein concentration

The reaction was measured at 240 nm for 3 min and H_2O_2 consumption was calculated using extinction coefficient, 39.4 mM^{-1} cm^{-1}.

3.9 Haematological Analysis

The whole blood of sacrificed animals was collected into EDTA bottles and analyzed for haematological parameters such as hemoglobin concentration (Hb), white blood cell count (WBC), red blood cell count (RBC), packed cell volume (PCV), platelet (PLT), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular

haemoglobin concentration (MCHC). The determination of hematological parameters was done within two hours of sample collection using Automated Hematological analyzer.

3.10 Histological Analysis of the Colon and other Organs

The colon tissue was rapidly fixed for 18-24 hours in 10 % formal saline and dehydrated in ascending grades of alcohol (70%, 80%, 90%, 95%, and 100 % respectively) for 2 hours each. The tissue was cleared in xylene and subsequently transferred through a pure paraffin in an oven for 1 hour and then into a second pot of melted paraffin for additional 2 hours (infiltration). The tissues were immersed in a mould containing molten paraffin wax, which is allowed to solidify (embedding) while the tissue is inside and ready for sectioning. Sections of 3 µm in thickness were prepared according to standard micro techniques onto glass slide and stained with hematoxylin and eosin. Photomicrographs under light microscope at high power magnifications (× 250) were examined for histopathological changes.

3.11 Carcinoembryonic Antigen Assay

The concentration of CEA was determined by enzyme-linked immunosorbent assay (ELISA) test which was based on the principle of a solid phase enzyme-linked immunosorbent assay. The assay system utilizes a monoclonal antibody directed against a distinct antigenic determinant on the intact CEA molecule was used for solid phase immobilization (on the microtiter wells). A goat anti-CEA antibody conjugated to horseradish peroxidase (HRP) is in the antibody-enzyme conjugate solution. The test sample was allowed to react simultaneously with the two antibodies, resulting in the CEA molecules being sandwiched between the solid phase and enzyme-linked antibodies. After 1 hour incubation at room temperature, the wells were washed with water to remove unbound labelled antibodies. The color development was stopped with the addition of Stop Solution changing the color to yellow. The concentration of CEA is directly proportional to

the color intensity of the test sample. Absorbance was measured spectrophotometrically at 450 nm.

3.12 Immunohistochemical Analysis

This analysis is based on the principle that the any damaged DNA mismatch repair gene(MLH1) by the MNU carcinogen present in the tissue samples will serve as an antigen that binds to the MLH1 antibody to form a complex. This complex subsequently produces a color that can be seen using microscope. The method used is the Avidin Biotin Complex (ABC) method also referred to as the avidin biotin Immunoperoxidase method, Gold *et al* (1965). Exactly 2 microns thick of formalin fixed and paraffin embedded tissues were cut for the IHC. Antigenic sites were retrieved using citric acid (pH 6.0) solution in pressure cooker at 100^0C for 25 minutes. Peroxidases, protein and biotin blocks were done using hydrogen peroxide, avidine and biotin respectively. *MLH1* antibody was diluted (1.100) and incubated on sections at 60^0C for 60 minutes. These were followed by the biotylinated secondary antibody, streptavidine and DAB/substrate reaction as described by Goldstein *et al*., (2007); Wolff *et al*., (2007).

3.13 Statistical Analysis

Data was analyzed using SPSS version 20.0. The statistical significance between the control and other groups of experimental animals was determined by one-way ANOVA followed by Duncan test for multiple comparisons. Statistical test was performed at $p \leq 0.05$ level of significance, Day *et al* (1989). The results were set as Mean ± SD.

CHAPTER FOUR

4.0 RESULTS

Cucurbita maxima fruit inclusion on packed cell volume (PCV), neutrophil, lymphocyte revealed that the MNU control group showed a lower percentage level of packed cell volume (PCV) which was significantly ($p \leq 0.05$) different from the normal control group and the groups that received various percentage of dietary *Cucurbita maxima* fruit inclusion. The MNU control group showed higher percentage level of neutrophil and lymphocyte density which was significantly ($p \leq 0.05$) difference from the normal control group and the group on different percentages of dietary *Cucurbita maxima* fruit inclusion (2.5%, 5% and 10%) in both colon and liver respectively (Figure 4.1).

Effect of dietary *Cucurbita maxima* fruit inclusion on haemoglobin and platelet concentration in the blood serum of different experimental group revealed that the MNU control group showed lower level of Haemoglobin and Platelet concentration which was significantly ($p \leq 0.05$) different from the normal control group and the group on different percentages of dietary *Cucurbita maxima* fruit inclusion (2.5%, 5% and 10%) in both colon and liver respectively (Figure 4.2).




Figure 4.1: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Packed Cell Volume, Neutrophil and Lymphocyte Values following 12 Weeks MNU Administered in Different Experimental Groups. Values are Mean±SD. Values having different letters across the column are significantly (p<0.05) different




Figure 4.2: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Haemoglobin and Platelet Concentration following 12 Weeks MNU Administered in Different Experimental Groups. Values are Mean±SD

Effect of dietary *Cucurbita maxima* fruit inclusion on white blood cell and red blood cell counts in the blood of different experimental group revealed that the MNU control group showed higher level of white blood cell counts which was significant ($p \leq 0.05$) difference from the normal control group and the group that received various percentage of dietary *Cucurbita maxima* fruit inclusion. The MNU control group showed lower level of red blood cell counts which was significant ($p \leq 0.05$) difference from the normal control group and the group on different

percentages of dietary *Cucurbita maxima* fruit inclusion (2.5%, 5% and 10%) in both colon and liver respectively (Figure 4.3)




Figure 4.3: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on White Blood Cell and Red Blood Cell following 12 Weeks MNU Administered in Different Experimental Groups

Values are Mean±SD

Table 4.1 displays the result of Oxidative Stress-related Biochemical Parameters effects on Lipid Peroxidation- LPO. There is a significantly increase in MDA level of the MNU treated control group which was significantly ($p \leq 0.05$) higher than the normal control group and other groups on different percentages of dietary *Cucurbita maxima* fruit inclusion (2.5%, 5% and 10%) in both colon and liver respectively.

Assessment of the antioxidant enzyme catalase in the liver and colon homogenate, revealed a lower activity for MNU control group which was significant differences ($p \leq 0.05$) from the normal control group the other groups on different percentages of dietary *Cucurbita maxima* fruit inclusion (2.5%, 5% and 10%) in both colon and liver respectively (Table 4.2).

Assessment of the antioxidant enzyme Superoxide dismutase in the liver and colon homogenate, revealed a lower activity for MNU control group which was significant differences ($p \leq 0.05$) from the normal control group the other groups on different percentages of dietary *Cucurbita maxima* fruit inclusion (2.5%, 5% and 10%) in both colon and liver respectively (Table 4.3).

Table 4.1: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Melon aldehyde levels following 12 Weeks Intrarectal MNU Instillation

GROUP TREATMENT	MDA levels (nmol /mg protein)	
	Colon	liver
A Normal Control	**113.50±0.9^{c}**	**56.00±4.4^{c}**
B MNU Control	**236.81±10.7^{d}**	**96.97±1.5^{d}**
C MNU+2.5% of *C. maxima* fruit	**116.73±1.1^{c}**	**51.96±3.9^{c}**
D MNU+5% of *C. maxima* fruit	**60.25±2.6^{b}**	**29.59±3.2^{b}**
E MNU+10% of *C. maxima* fruit	**28.59±1.22^{a}**	**28.51±2.8^{b}**
F 10% Dietary Control	**20.73±1.6^{a}**	**16.69±1.4^{a}**

Values are Means±SD; Values having different letters down the column are significantly different ($p \leq 0.05$)

Table 4.2: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Catalase (CAT) Activity following 12 Weeks Intrarectal MNU Instillation

GROUP TREATMENT	CAT levels (U/mg protein)	
	Colon	liver
A Normal Control	29.92±0.9^{d}	54.57±7.59^{c}
B MNU Control	11.50±1.8.0^{a}	31.76±3.4^{a}
C MNU+2.5% of *C. maxima* fruit	16.76±3.7^{ab}	41.51±2.2^{b}
D MNU+5% of *C. maxima* fruit	21.38±2.0^{bc}	53.89±2.7^{c}
E MNU+10% of *C. maxima* fruit	27.97±1.3^{cd}	62.36±2.1^{c}
F 10% Dieatry Control	37.63±10.3^{e}	71.38±6.8^{d}

Values are Means±SD; Values having different letters down the column are significantly ($p \leq 0.05$) different

Table 4.3: Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Superoxide dismutase Activity following 12 Weeks Intrarectal MNU Instillation

GROUP TREATMENT	SOD levels (U/ml/mg protein)	
	Colon	liver
A Normal Control	**264.37±1.8^{c}**	**69.51±1.0^{b}**
B MNU Control	**212.89±4.5^{a}**	**68.44±6.4^{a}**
C MNU+2.5% of *C. maxima* fruit	**240.05±7.2^{b}**	**68.31±2.6^{b}**
D MNU+5% of *C. maxima* fruit	**292.81±12.0^{d}**	**73.78±4.2^{b}**
E MNU+10% of *C. maxima* fruit	**316.67±3.7^{e}**	**81.89±1.2^{c}**
F 10% Dietary Control	**275.48±3.5^{e}**	**88.21±0.8^{d}**

Values are Means±SD; Values having different letters down the column are significantly different ($p \leq 0.05$)


The Photomicrograph of histological section of the colon following 12 weeks MNU administration in rats showing essentially normal colon in the normal control group and the

groups that are treated with 10% of dietary *Cucurbita maxima* fruit inclusion with MNU and normal saline respectively, but there is focal inflammation in the MNU control group and the groups treated with 2.5% and 5% dietary *Cucurbita maxima* fruit inclusion (Plate 4.1).


The Photomicrograph of histological section of the liver following 12 weeks of MNU administration in rats showed essentially normal liver in the normal control group and the groups that are treated with higher concentration of MNU (5% and 10%) of dietary *Cucurbita maxima* fruit inclusion with MNU and normal saline respectively, but there is focal inflammation in the MNU control group and the group treated with 2.5% dietary *Cucurbita maxima* fruit inclusion (Plate 4.2).

Effect of dietary *Cucurbita maxima* fruit inclusion on carcinoembryonic antigen (CEA) in the blood serum of different experimental group revealed that the MNU control group showed higher level of CEA concentration which was significantly ($p \leq 0.05$) different from the normal control group and the group that received 2.5%, 5% and 10% of dietary *Cucurbita maxima* fruit inclusion (Figure 4.4).


The Photomicrograph of immunohistochemistry section of the colon following 12 weeks MNU administration in rats showed that there is moderate to mild expression of MLH1 antibody in the normal control group and the groups treated with 10% dietary *Cucurbita maxima* fruit inclusion, but there is no expression of MLH1 antibody in the MNU control group and the groups that received 2.5% and 5% of dietary *Cucurbita maxima* fruit inclusion (plate 4.3).




1a: Normal control group showing essentially normal colon. (X40).




1b: MNU control group showing essentially normal colon but with mild (a)interglandular inflamation. (X40).




1c: MNU + 2.5% showing essentially normal colon but with (a) deeper mucosal inflammation and (b) mild ulceration. (x40)



1d: MNU + 5% showing essentially normal colon but with (a) moderate mucosal inflammation. (x40)




1e: Normal control group showing essentially normal colon. (X40).




1f: Normal control group showing essentially normal colon. (X40).


Plates 4.1: Photomicrogragh of Histopathological Section of the Colon following 12 Weeks MNU Administration in Rats Fed with different Level of Dietary *Cucurbita maxima* Fruit Inclusion Diet




2a: Normal control group showing essentially normal liver (X40).




2b: Normal control group showing essentially normal colon. (X40).




2c: Normal control group showing essentially normal colon. (X40).



2d: Normal control group showing essentially normal colon. (X40).



2e: Normal control group showing essentially normal colon. (X40).



2f: Normal control group showing essentially normal colon. (X40).

Plates 4.2: Photomicrograph of Histopathological Section of the Liver following 12 Weeks MNU Administration in Rats Fed with different Level of Dietary *Cucurbita maxima* Fruit Inclusion Diet.







Figure 4.7: Carcinoembryonic antigen Concentration of the Serum following Intrarectal Administration of MNU in Rats Fed with Different Levels of Dietary *C. maxima* Fruit Inclusion.




3a: Normal control group showing no expression of the antibody (MLH1).




3b: MNU control group showing moderate expression of the antibody (MLH1). (x40)




3c: MNU+2.5 showing moderate expression of the antibody (MLH1). (x40)



3d: MNU+5% showing mild expression of the antibody (MLH1). (x40)



3e: MNU+10% showing mild expression of the antibody (MLH1). (x40)



3f: Dietary control group showing no expression of the antibody (MLH1). (x40)

Plates 4.3: Photomicrogragh of Immunohistochemistry Section of the Colon, following 12 Weeks MNU Administration in Rats Fed with Different Level of Dietary *C. maxima* Fruit Inclusion.

CHAPTER FIVE

5.0 DISCUSSION

Dietary inclusion of *Cucurbita maxima* fruit showed result of hematological indices of rats fed with various diet mixtures. The significant ($P \leq 0.05$) decrease in the packed cell volume (PCV) and hemoglobin (Hb) concentration of the MNU control group indicating anemia, which may be due to damage to red blood cell, Mani *et al.,* (2014), which in this research is connected with MNU toxicity. Thus, low PCV level might suggest oxidative stress-related RBC damage in the MNU treated control group, while the increase level of PCV in the *Cucurbita maxima* powdered treated groups with percentage inclusion of 2.5%, 5% and 10%, suggest the potency of the dietary *Cucurbita maxima* fruit inclusion with its phenolic contents, Sopan *et al.,* (2014) to alleviate oxidative stress-related damage to red blood cell membrane. The significant ($p \leq 0.05$) increase in WBC in MNU control group compared to the normal control group which is not significantly ($p \geq 0.05$) different from other groups that received 2.5% ,5% and 10% dietary inclusion of *Cucurbita maxima,* suggested that the plant has potency in boosting the immune system, Sapon *et al.,* (2014) (Figure 4.3).

Oxidative stress has been suggested to contribute to the pathogenesis of carcinogenicity and lipid peroxidation which is one of the characteristic features of cancer, Halliwell (1994). The reactive intermediates, produced by oxidative stress, can alter the membrane bilayers and cause the lipid peroxidation of polyunsaturated fatty acids (PUFA) leading to the formation of lipoperoxyl radical (LOO^{-}) which in turn, reacts with the lipid to yield a lipid radical and a lipid hydroperoxide (LOOH). Lipid hydroperoxide LOOHs are unstable and they generate new

peroxyl and alkoxyl radicals and decompose into secondary products, Halliwel and Chirico, (1993). Malondialdehyde (MDA) is found during oxidative degeneration as a product of free oxygen radicals which is accepted as an indicator of lipid peroxidation MDA, the end product of lipid peroxidation, was reported to be higher in cancer tissues than in non-diseased organ, Nielsen *et al.*, (1994).

In the present study, N-methyl-N-nitrosourea (MNU) intoxication resulted in lipid peroxidation (LPO), which produced MDA as by product that caused damage to the cell membranes (Moller *et al.,* 1998; Siddhartha *et al.,* 2011). The groups that were fed various percentages of *Cucurbita maxima* fruit powder (2.5%, 5% and 10%) indicated significant decrease levels of MDA in the Colon and Liver respectively which suggest that the lipid peroxidation of membranes caused by MNU intoxication was prevented by inclusion of the *Cucurbita maxima* fruit powder and thus implying that this fruit contain antioxidant compounds. This antioxidant activity could be due to compounds such as phenolic acid and phenolic diterpenes, Sapon *et al.,* (2014), that were capable of scavenging free radicals from the tissues of the treated rats.

According to Gacche *et al*., (2010), 2,2–diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity of *Cucurbita maxima* fruit is 41%, inhibition of LPO is 40.3%, Vitamin C is 15.36mg, Total phenols is 13.30mg and FICA 22.3% respectively. A redox balance between pro-oxidants and antioxidants is essential for the normal cellular functioning, Nordberg and Amer, (2001); Rahal *et al.,* (2014). Impairment in the ratio of oxidants and antioxidants initiates in the patho-physiological events that culminate in molecular and cellular damage to macromolecules and vital organs, Valko *et al*., (2006).

The antioxidants SOD and CAT play an important role in reducing cellular stress. SOD scavenges the superoxide radical by converting it to hydrogen peroxide and molecular oxygen, Robinson (1998), while CAT brings about the reduction of hydrogen peroxide and protects higher tissues from highly reactive hydroxyl radicals Brioukhanov and Netrusov, (2004).

In the present study, the activity of CAT was significantly depleted in MNU control group rats. Dietary *Cucurbita maxima* fruit inclusion significantly ($p \leq 0.05$) increased the CAT activity in the group that received various percentages of 2.5%, 5% and 10% and also in the 10% dietary control group rats when compared to MNU control group rats. The SOD activity was significantly decreased in rats exposed to only MNU, dietary *Cucurbita maxima* fruit inclusion significantly ($p \leq 0.05$) increased the SOD activity in the groups which took the 2.5%, 5% and 10% respectively and also in 10% inclusion control group. Dietary *Cucurbita maxima* fruit has excellent antioxidant activity which may protect individuals from oxygen free radicals and lipid peroxidation induced damage, Rahal *et al.,* (2014)

According to Thompson and Adlakha (1991), MNU rat model have several advantages such as reliability of tumour induction organs site specificity, tumour of ductal origin and predominantly carcinomatous histopathologic characterization and the ability to examine tumour initiation and promotion processes. The histopathological findings of the colon and liver following 12 weeks MNU administration in rats in (Plate 4.1 and Plate 4.2) showing essentially normal colon and liver in the normal control group and the groups that were treated with 5% and 10% dietary inclusion of *Cucurbita maxima* fruit, strongly suggest high anti-oxidative effect of the *Cucurbita maxima* fruit, which may be as a result of it high contents of antioxidant compounds such as polyphenols and antioxidant vitamins, Muntean *et al.,* (2013). Inflammation observed in the

MNU control group and the groups treated with 2.5% dietary inclusion of *Cucurbita maxima* fruit, revealed that MNU caused severe damage to the .mucosal cells, which is connected with MNU toxicity and that higher percent inclusion is necessary for significant effect.

Immunohistochemistry has been widely used in diagnosing carcinogenesis because it takes advantage of the principle of the antigen-antibody affinity reaction through its ability to identify and localize proteins of interest via detection with labelled conjugates Mann (1902). These antigens produced in disease condition have corresponding antibodies produced by the body to form a complex compounds which are usually picked up by immunochemical dyes or stains giving a distinctive colour (brownish colour in the case of the *Mut L homoloq* 1 gene) as "expression" under the microscope. The distinct moderate to mild expression of *Mut L-homoloq* 1 antibody in the normal control group and the groups treated with 5% and 10% dietary inclusion of *Cucurbita maxima* fruit in (Plate 4.3) strongly suggest high anti-oxidative effect of the *Cucurbita maxima* fruit, Sapon *et al.,* (2014), which may be as a result of its high contents of antioxidant compounds including polyphenols and antioxidant vitamins. In contrast there was expression of *Mut L homoloq* 1 antibody in the MNU control group and the groups treated with 2.5% and 5% dietary inclusion of *Cucurbita maxima* fruit, due to damaged or mutated *Mut L-homoloq* 1 gene Sapon *et al.*, (2014). According to Thompson *et al*. (1995) and Welsch *et al*., (1985), defect in the DNA mismatch repair gene like the *Mut L homoloq* 1 gene can lead to its inability to recognize and repair errors that occur during DNA replication. This loss of DNA mismatch repair gene activity accelerates the rate of accumulation of mutations in other gene leading to carcinogenesis, Sapon *et al.*, (2014); Kim *et al.,* (2004).

CEA is a glycoprotein which is normally present at only very low concentrations in adult plasma but its concentration can be increased during the process of carcinogenesis or in the presences of tumors particularly colorectal cancers, Herbert *et al.,* (1980). They are well known biomarker use to determine the extent of preoperative colon cancer, Hajime *et al.,* (2012). The significant ($p \leq 0.05$) increase in the level of CEA concentration of MNU control group may be as a result of MNU toxicity. According to Sukumar *et al.,* (1983) MNU toxicity can lead to inflammation and damage to normal mucosal cells, while a significant ($p \leq 0.05$) decrease in the level of CEA concentration in the groups treated with 2.5%, 5% and 10% of dietary *Cucurbita maxima* fruit inclusion strongly suggests the potency of the extract to prevent oxidative-stress related damage to cell membrane do to their ability to absorb, neutralize and to quench free radicals, which may be as a result of it high contents of antioxidant compounds such as polyphenols and flavonoids, Muntean *et al.,* (2013).

CHAPTER SIX

6.1 SUMMARY

This study established that;

(i) *Cucurbita maxima* fruit significantly ($P \leq 0.05$) protected haematological indices (Packed cell volume, Haemoglobin, Red blood cell and White blood cell from MNU-induced toxicity.

(ii) *Cucurbita maxima* fruit significantly ($P \leq 0.05$) prevented protein damage in the organs (liver and colon) from MNU-induced damage and damage to the DNA mismatch repair gene (*Mut L homoloq* 1) by the MNU carcinogen.

(iii) *Cucurbita maxima* fruit prevented oxidative stress in colon carcinogenesis by increasing the activities of catalase and SOD, and also by decreasing the concentration of MDA in the organs (liver and colon).

6.2 CONCLUSION

Cucurbita maxima fruit has potential to prevent colon carcinogenesis and protect against MNU-induced oxidative stress, organs damage and its adverse effects on haematological parameters.

6.3 RECOMMENDATION

From this study it is recommended that people should be encouraged to increase their consumption of *Cucurbita maxima* fruit because it has high antioxidants which play vital roles in prevention of cancer and other diseases.

REFERENCES

Adams G.G., Imran S., Wang S., Mohammad A., Kok S., Gray D.A., Channell G.A., Morris G.A., Harding S.E., 2011. The hypoglycaemic effect of pumpkins as anti- diabetic and functional medicines. Food Res. Internation., 44: 862-867.

Afsar U.A. (2014). *An overview of inflammation: mechanism and consequences*. Front. Biol. 2011, 6(4): 274–281

Ager A, May MJ. Understanding high endothelial venules: Lessons for cancer immunology. Oncoimmunology. 2015;4(6):e1008791.

Aggarwal B B, Vijayalekshmi R V, Sung B (2009). Targeting inflammatory pathways for prevention and therapy of cancer: short term friend, long-term foe. Clin Cancer Res, 15(2): 425–430

Ahmed AA. The surgical management and outcome of oesophageal cancer in Addis Ababa. Ethiop Med J. 2000; 38: 147–152.

American Cancer Society, Facts and figures. 1999

Anderson R L, Wolf W J. (1995). Compositional changes in trypsin inhibitors, phytic acid, saponins and isoflavones related to soybean processing. J. Nutr. 125: 581S-588S.

Anuj Y, Rewa K, Ashwani Y, J.P. Mishra, Seweta S and Shashi P (2016). *Antioxidants and its functions in human body - A Review.* Res. Environ. Life Sci. **9**(11) 1328-1331 https://www.researchgate.net/publication/311674771

AOAC, 2001. Official methods of analysis. Association of official analytical chemists arlington, VA. 2001.

Armstrong, D., and Doll, R.: Environmental factors and cancer incidence and mortality i different countries, with special reference to dietary practices. Int. J. Cancer 15:6 17-63 1, 1975

Aslani B.A. and Ghobadi S. (2016). Studies on oxidants and antioxidants with a brief glance at their relevance to the immune system. Life Sci. 146: 163–173.

Attarde DL, Kadu SS, Chaudhari BJ, Kale SS, Bhambre RS. In vitro Antioxidant activity of pericarp of Cucurbita maxima Duch. Ex Lam. International journal of Pharm Tech Research 2010; 2(2):1533-1538.

Avrameas, S., and Ternynck, T. (1971). Peroxidase labelled antibody and Fab conjugates with enhanced intracellular penetration. *Immunohistochemistry,* **8,** 1175-1179

Aderemi, F. A. (2004). Effects of replacement of wheat bran with cassava root sieviate supplemented or unsupplemented with enzyme on the haematology and serum biochemistry of pullet chicks. *Tropical Journal of Animal Science, 7,* 147-153.

Adebayo, O. R, Forombi A. G & Oyekanmi A. M. (2013). Proximate, Mineral and Anti-nutritonal evaluation of pumpkin pulp (Cucurbita pepo). Journal of applied chemistry. 4(5): 25-28.

Aebi, H.E, (1983). Catalase. In: Methods of Enzymatic Analysis, Bergmeyer, H.U. (Ed). 3rd Edn., Verlag Chemie Weinhem, pp: 276-286.

Afolabi, K. D., Akinsoyinu, A. O., Olajide, R., and Akinleye, S. B. (2010). Haematological parameters of the Nigerian local grower chickens fed varying dietary levels of palm kernel cake. *Proceedings of 35th Annual Conference of Nigerian Society for Animal Production (p.247).*

Aster, J. C. (2004). Anaemia of diminished erythropoiesis. In V. Kumar, A. K. Abbas, N. Fausto, S. L. Robbins, & R. S. Cotran (Eds.), Robbins and Cotran Pathologic Basis of Disease *Saunders Co. Philadelphia*. 7th ed., p.638-649.

Atawodi, S.E., 2010. Polyphenol Composition and in vitro Antioxidant Potential of Nigerian *Canarium schweinfurthii* English Oil. *Advance Biological Research,* 4: 314-322.

Atawodi, S.E., 2011. Polyphenol Content and *In Vitro* Antioxidant Activityof methanol extract of Seeds of *Irvingia gabonensis* Baill of Nigerian Origin. *Elect. Journal Environment Agriculture Food Chemistry,* 10:2314-2321.

Atawodi, S.E., (2011). Nigerian foodstuffs with prostate cancer chemopreventive polyphenols. Proceedings, science of global prostate cancer disparities in black men conference. *Infectious Agents and Cancer,* 6: S2-S9.

Atawodi, S.E., (2011). Evaluation of the hypoglycemic, hypolipidemic andantioxidant effects of methanolic extract of "Ata-ofa" polyherbal tea(*A-Polyherbal*) in alloxan-induced diabetic rats. *Drug Invention Today*,3: 270-276.

Atawodi, S.E., Atawodi, J.C. Idakwo, G.A. Pfundstein, B. Haubner, R. Wurtele, G. Spiegelhalder, B. Bartsch, H. and Owen, R.W. (2009a). Polyphenol composition and antioxidant potential of *Hibiscusesculentus* L. Fruit Cultivated in Nigeria. *J. Medline Food,* 12:1316-1320.

Attarde D.L., Kadu S.S., Chaudhari B.J., Kale S.S., Bhamber R.S., (2010). In vitro antioxidant activity of pericarp of *Cucurbita maxima* Duch. Ex Lam. *International Journal of Pharmatech Research*, 2, 1533-1538.

Atawodi, S., Ayo, J. O., Atodo, A. D., & Dzende, T. (2005). Some haematological parameters and the erythrocyte osomotic fragility in the laughing dove (Streptopella senegalensis) and the village weaner bird (Ploceus cucullatus*). Proceedings of the 10th Annual Conference of Animal Science Association of Nigeria.* (p.384-387).

Barber, D. A., and Harris, S. R. (1994). Oxygen free radicals and antioxidants: A review. Am Pharm NS34, 26–35.

Bergomas F, Grizzi F, Doni A, Pesce S, Laghi L, Allavena P, et al. Tertiary intratumor lymphoid tissue in colo-rectal cancer. Cancers (Basel). 2011;4(1):1-10.

Balaguer, F.; Link, A.; Lozano, J. J.; Cuatrecasas, M.; Nagasaka,T.; Boland, C. R.; Goel, A. (2010). "Epigenetic Silencing of miR-137 is an Early Event in Colorectal Carcinogenesis". *Cancer Research*. **70** (16): 6609–6618.

Bamishaiye, E. I., Muhammad, N. O., & Bamishaiye, O. M. (2009). Haematological parameters of albino rats fed on tiger nuts (Cyperus esculentus) tuber oil meal-based diet. *The International Journal of Nutrition and Wellness, 10*(1).

Bernstein, Carol; Holubec, Hana; Bhattacharyya, Achyut K.; Nguyen, Huy; Payne, Claire M.; Zaitlin, Beryl; Bernstein,Harris (2011). "Carcinogenicity of deoxycholate, a secondary bile acid". *Archives of Toxicology*. **85** (8): 863–71.

Boni C., Andre T., Navarro M., Tabernero J., Hickish T., Topham C., Bonetti A., Clingan P., Bridgewater J., Rivera F., and Aimery de Gramont J. (2009). Improved Overall Survival With Oxaliplatin, Fluorouracil, and Leucovorin As Adjuvant Treatment in Stage II or III Colon Cancer in the MOSAIC Trial. *Clinical Oncology* 27: 309-36.

Boucher D, Cournoyer D, Stanners CP, fuks A. Studies on the control of gene expression of the carcinoembryonic antigen family in human tissue. Cancer Res 1989;49:847-52

Bose KS, Vyas P, Singh M (2012). Plasma non-enzymatic antioxidants Vitamin C, E, beta-carotene, reduced glutathione levels and total antioxidant activity in Oral sub mucous fibrosis. Eur Rev med pharmacol Sci, Oxford, UK.

Boxin, O.U., Dejian H., Maureen A.F. and Elizabeth K.D.: Analysis of antioxidant activities of common vegetables em(ploying oxygen radical Absorbance Capacity (ORAC) and Ferric Reducing Antioxidant Power (FRAP) Assays: A comparative study. J. Agric. Food Chem., 5: 223-228 (2002)

Boyle T, Keegel T, Bull F, Heyworth J, Fritschi L. (2012). Physical activity and risks of proximal and distal colon cancers: a systematic review and meta-analysis. *Journal of National Cancer Institute*. 104: 1548-1561.

Brahmachari, U.N. (2001). Herbal Drugs. Current Science; 81:15-16.

Brunken, U., Schmidt, M., Dressler, S., Janssen, T., Thombiano, A. & Zizka, G. (2008). West African plants - A Photo Guide. Forschungsinstitut Senckenberg, Frankfurt/Main.

Cadet, J.L. and Kahler, L.A. (1994). Freeradical mechanisms in schizophrenia andtardive dyskynesia. *Neurological Science Biobehaviour. Research.*18 (4), 457-467.

Chaudiere J. and Frerrari-LLiou R. (1999). Intracellular antioxidants: from chemical to biochemical mechanisms. Food Chem Toxicol. 37: 949–962.

Cancer Research UK 2011 April 2011.

Carter P. (1971). Spectrophotometric determination of serum iron at the submicrogram level with a new reagent (ferrozine). *Analytical Biochemistry*, 40: 450–455.

Carter. J. (1993).Dietary fiber guide. Cereal Foods World, 38 (10): 755-759.

Chineke, C. A., Ologun, A. G., and Ikeobi, C. O. N. (2006). Haematological parameters in rabbit breeds and crosses in humid tropics. *Pakistan Journal of Biological Sciences, 9*(11), 2102-2106.

Colorectal cancer: The diagnosis and management of colorectal cancer; NICE Clinical Guideline (November 2011)

Coppola D, Nebozhyn M, Khalil F, Dai H, Yeatman T, Loboda A, et al. Unique ectopic lymph-node-like structures present in human primary colorectal carcinoma are identified by immune gene array profiling. The American journal of pathology. 2011;179(1):37-45.

Cunningham JM, Kim CY, Christensen ER, Tester DJ, Parc Y, Burgart LJ, et al. The frequency of hereditary defective mismatch repair in a prospective series of unselected colorectal carcinomas. Am J Hum Genet. 2011; 69: 780-90.

Cunningham JM, Christensen ER, Tester DJ, Kim CY, Roche PC, Burgart LJ, et al. Hypermethylation of the hMLH1 promoter in colon cancer with microsatellite instability. Cancer Res. 2001; 58: 3455-60.

Croce, C.M. (2008). "Oncogenes and cancer". *The New England Journal of Medicine*. 358 (5): 502–11.

Croshaw, R., Willis, J., Dawson, D., Shipitsin, M., Willson, J.K., Sukumar, S., Polyak, K., Park, B.H., Pethiyagoda, C.L., Pant, P.V., Ballinger, D.G., Sparks, A.B., Hartigan, J., Smith, D.R., Suh, E., Papadopoulos, N., Buckhaults, P., Markowitz, S.D., Parmigiani, G., Kinzler, K.W., Velculescu, V.E., Vogelstein, B. (2007). "The genomic landscapes of human breast and colorectal cancers". *Science*. 318 (5853): 1108–1113. doi:10.1126/science.1145720. PMID 17932254.

Daramola, J. O., Adeloye, A. A., Fatoba, T. A., and Soladoye, A. O. (2005).Haematological and biochemical parameters of West African Dwarf goats. *Livestock Research for Rural Development, 17*(8), 95. Retrieved January 22, 2014, from http://www.lrrd.org/lrrd17/8/dara17095.htm.

Das R., Sahu J., Gupta R.N., (2010), Potentiative Activity of Cucurbita Maxima Seed Extract With Beta Vulgaris & Smilax Regelii Root Extract To Reduce Extra Fats From The Body. International Journal of Pharmaceutical Sciences and Research, 1, p.57-62.

Das, R. et al. Potentiative Activity Of Cucurbita Maxima Seed Extract With Beta Vulgaris & Smilax Regelii Root Extract To Reduce Extra Fats From The Body. *International Journal of Pharmaceutical Sciences and Research* 2010; 1:57-62. 10

Day. R. W., and G.P. Quinn (1989) "comparison of treatments after an analysis of variance in ecology." *Ecological Monographs* 59:433-463

Davidson LA, Jiang YH, Derr JN, Aukema HM, Lupton JR, Chapkin RS. Protein kinase C isoforms in human and rat colonic mucosa. Arch Biochem Biophys 1994;312:547–53.

de Martel C, Franceschi S (2009). Infections and cancer: established associations and new hypotheses. Crit Rev Oncol Hematol, 70(3): 183–194

De Martel C, Ferlay J, Franceschi S, et al. Global burden of cancers attributable to infections in 2008: a review and synthetic analysis. Lancet Oncol 2012; 13:607-15.

Demple, B., and Harrison, L. (1994). Repair of oxidative damage to DNA: Enzymology and biology. Annu Rev Biochem 63, 915–48.

Dean PN, Jett JH. (1974). Brief note: mathematical analysis of DNA distributions derived from flow microfluorometry. J Cell Biol. 60: 523.

Dizdaroglu, M. (1992). Oxidative damage to DNA in mammalian chromatin. Mutat Res 275, 331–42

Duduyemi BM, Oluwasola AO, Akang EEU, Thomas-Ogunniyi JO. A 16-year review of clinico-pathological pattern of colorectal carcinoma at University College Hospital, Ibadan. Nigerian Journal of Gastroenterology and Hepatology Vol.3 Nos. 1&2 June & Dec., 2013.

DuffyMJ; Serum tumor markers in breast cancer: are theyof clinical value? Clin Chem. 2006 Mar;52(3):345-51. Epub 2006 Jan 12

Dow LE, O'rourke KP, Simon J, Tschaharganeh DF, Van Es JH, Clevers H, et al. Apc restoration promotes cellular differentiation and reestablishes crypt homeostasis in colorectal cancer. Cell. 2015; 161: 1539-52

Doren, E.T. (1997). Vegetable ivory and other palm nuts/seeds as an art/craft medium. *Journal of the International Palm Society*. (renamed as Palm since 1999), 41 (4), 18-25.

Dreosti I.E., (1991). Trace elements, micronutrients, and free radicals. Totowa: *Human Press.* pp. 149-199.

Dusica, P. and Vesna, T. (2002) Oxidative stress as marker of positive symptoms in schizophrenia. Facta Universitatis. *Medicine & Biology*. 9 (2), 157-161.

Dwivedi, V. Shrivastava, R. Hassan, S. Ganguly, C. Bharadwaj, M. Ganguly, C. (2011) :Flavoring agents used in Indian cooking and their anticarcinogenic properties. *Asian Pacific Journal of Cancer Prevention*, 11, 25-28.

El-Gendy, A.; El-Mileegy, A.; Ghyaty, E.; Malek, H.; El-Hamid Mousa, A. (2008). "The Beneficial Dietary Hypotensive and Hypolipidaemic Effects of Hyphaene Thebaica (Doum)". *Internet Journal of Alternative Medicine 7 (1).*

Fakih MG, Padmanabhan A;CEAmonitoring in colorectal cancer. What you should know.Oncology(Williston Park). 2006 May;20(6):579-87; discussion 588, 594, 596 passim.

Fraga, C. G., Shigenaga, M. K., Park, J. W., Degan, P., and Ames, B. N. (1990). Oxidative damage to DNA during aging: 8-hydroxy-2'-deoxyguanosine in rat organ DNA and urine. Proc Natl Acad Sci U S A 87, 4533–7.

Fanburg BL, Lee SL (2015). A new role for an old molecule: serotonin as a mitogen. Am J physiol 272: L795-L806

FAOSTAT (Food and Agriculture Organization of the United Nation) 2013. http://faostat.fao.org /site/567/ Desktop Default. aspx? Page ID=567#ancor.

Fearon ER, Vogelstein B. A genetic model for colorectal tumorigenesis. Cell 1990;61:759–67.

Fernanda L K, Danielle C G M, Lívia C AR, Marisa C PP, Graciela FV, Lucas LC, Elizeu AR Iracilda ZC. (2012). A soy-based product fermented by Enterococcus faecium and Lactobacillus helveticus inhibits the development of murine breast adenocarcinoma. Food Chem Toxicol. 50: 4144–4148.

Fernandez Z.F., Guerra N.B., Diniz N.M., Salgado S., Guerra T.M., Lopez A.C., Neta J.C., Padilha M.R. (1998). Development of a milk beverage based on pumpkin flakes. Arch. Latinoam.Nutr.,48(2), 175-178.

Fernandez-PoI, J. A., Bono, V. H., and Johnson, G. S. Control of growth by picolinic acid: differential response of normal and trans formed cells. Proc. NatI. Acad. Sci. U. S. A., 74: 2889-2893, 1977.

Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray, F. GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC Cancer Base No. 11 [Internet]. Lyon, France: International Agency for Research on Cancer; 2013. Available from: http://globocan.iarc.fr, accessed on 18/12/2013.

Fischer-Nielsen, A., Jeding, I.B., Loft, S. (1994). Radiation-induced formation of 8-hydroxy-2'-deoxyguanosine and its prevention by scavengers. *Carcinogen.* 15, 1609-1612

Fletcher RH. Carcinoembryonic antigen. Ann Intern Med 1986;104:66-73

Friedman, D. L. Role of cyclic nucleotidesof cell growth and differen tiation. Physiol. Rev., 56: 652-708, 1976

Facciola, S., *Cornucopia, A Source Book of Edible Plants* Kampong.

FAOSTAT (2007). Agricultural Structure (Production, Price, Value). (available on-line at: http://apps.fao.org/faostat).

Faten, M. A. E. (2009). Antioxidant and anticancer activities of doum fruit extract (Hyphaenethebaica). *African Journal of Pure and Applied Chemistry*, 3 (10), 197-201.

Fearon, ER (1997). "Human cancer syndromes: Clues to the origin and nature of cancer". *Science*. 278 (5340):1043–1050. doi:10.1126/science.278.5340.1043. PMID 9353177.

Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray, F. GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet]. Lyon, France: *International Agency for Research on Cancer;* 2013. Available from: http://globocan.iarc.fr, accessed on 18/12/2016.

Ferlay, Soerjomataram I, Ervik M, Dikshit R, Eser S, et al (2012). Globocan, v 1.0, Cancer Incidence and Mortality Worldwide: IARC cancer Base No. 11 [Internet]. Lyon, Fance*: International agency for research on cancer:* 2013; Available from: http://globocan.iarc.fr.accessed on 13/12/2016.

Fletcher, R. (1997). Listing of useful plant of the world. Australian New crops.http://www.netcrops.uq.edu.au/listing/hyphaenethebaica.

Flynn, C. Edwards, R. and Kralj-Hans, I. (2007). Deoxycholic acid promotes the growth of colonic aberrant crypt foci. *Mol. Carcinog.*, 46, 60-70.

Gacche R.N, Kabaliye V.N, Dhole N.A, Jadhav A.D. (2010). Antioxidant potential of selected vegetables commonly used in diet in Asian subcontinent. *Journal of Natural Products and Resources* Vol.1(3), pp. 306-313

Garcia-Solis P, Alfaro Y, Anguiano B, Delgado G, Guzman RC, Nandi S, Diaz-Munoz M, Vazquez/Martinez O, Aceves C.(2005). Inhibition of N-methyl-N-nitrosourea induced mammary carcinogenesis by molecular iodine(I2) but not by iodide (I-) treatment Evidence that I2 prevents cancer promotion. Mol Cell Endocrinol. 236 49–57.

German JB (2013). Food processing and Lipid Oxidation. Adv Exp med Bio 459: 23-50

Ghanbari, R.F. Anwar, K.M. Alkharfy, A.H. Gilani and N. Saari, 2012. Valuable nutrients and functional bioactives in different parts of olive (*Olea europaea* L.)-A review. *International Journal Molecular Science*, 13: 3291-3340.

Globocan 2013.

Gold P, Freedman SO. Demonstration of tumor-specific antigens in human colonic carcinoma by immunological tolerance and absorption techniques

Gown AM. Genomic immunohistochemistry: a new era in diagnostic immunohistochemistry. Curr Diagn Pathol (2002); 8: 193-200.3.

Goodman, J., and Hochstein, P. Generation of free radicals and lipid peroxidation by redox cycling of Adriamycin and daunomycin. Bio chem. Biophys. Res. Commun., 77: 797-803, 1977.

Gregory, E. M., and Fridovich, I. Oxygen toxicity and the superoxide dismutase.J. Bacteriol., 114: 1193-1197, 1973.

Grivennikov S I, Greten F R, Karin M (2010). Immunity, inflammation, and cancer. Cell, 140(6): 883–899

Gutteridge JMC, Halliwel B (2011). Antioxidant in Nutrition, Health and Diseases. Oxford University Press, Oxford, UK

Gwanama C, Labuschagne MT, Botha AM (2000). Analysis of genetic variation in Cucurbita moschata by random amplified polymorphic DNA (RAPD) markers. Euphytica, 113: 19-24.

Halliwell, B. (1994). Free radicals and antioxidants: A personal view. *Nutr. Rev.* 52, 253-265.

Halliwell, B. Chiroco, S. (1993). Lipid peroxidation: its mechanism, measurement and significance. *Am. J. Clin. Nutr. 57, 715S-725S*

Halliwell, B. and Gutteridge, J.M.C.: The Definition and Measurement of Antioxidants in Biological Systems. Free Radical Biology and Medicinal., 18: 125-126 (1995).

Hammarstrom S. The carcinoembryonic antigen (CEA) family: structure, suggested functions and expression in normal and malignant tissue. Semin Cancer Biol 1999;9:67-81

Hernandez SM, Merrick CL, Eguilarte L (2005). Maintenance of squash (Cucurbita spp.) landrace diversity by farmers activities in Mexico. Genet. Resour. Crop Evol. 52: 697-707.

Herbert B, Bargrel A: A study of factors influencing plasma CEA levels in an unselected population. *Oncodev Biol Med* 1:191-198

Hostetter RB, Augustus LB, Mankarious R, Chi K, Fan D, Toth C, et al. carcinoembryonic antigen as a selective enhancer of colorectal cancer metastasis. J Natl Cancer Inst 1990;82:380-5

Jeffrey C. (2007). Key to the Cucurbitaceae of West Tropical Africa witha guide to localities of rare and little -known species. *Journal West.*

Jeffery M, Hickey BE, Hider PN; Follow-up strategies for patients treated for non-metastatic colorectal cancer.Cochrane Database SystRev. 2007 Jan 24;(1):CD002200.

Jessup JM, Thomas P. Carcinoembryonic antigen: function in metastasis by human colorectal carcinoma. Cancer Metastasis Rev 1989;8:263-80

Jose, M. A. et al. Diuretic activity of seeds of Cucurbita maxima duchesne in albino wistar rats. Indian J Pharmacol 2008; 40: S72

Johnson SM, Gulhati P, Rampy BA, Han Y, Rychahou PG, Doan HQ, et al. Novel expression patterns of PI3K/Akt/mTOR signaling pathway components in colorectal cancer. J Am Coll Surg. 2010; 210: 767-76, 776-8.

Jose M.A., Balasubramanian S., Rahman A.H., Varghese N.A., Kumar A.S., Sumaiya A., Vivekanandhan S., Sridharan C., Radhakrishnan M., (2008), Diuretic activity of seeds of *Cucurbita maxima* Duchesne in albino wistar rats. *Indian Journal Pharmacol*ogy. 40, 72**.**

Kabir, M., Akpa, G. N., Nwagu, B. I., Adeyinka, I. A., and Bello, U. I. (2011). Sexual dimorphism, breed and age characteristics of rabbits in Zaria, *Nigeria Proceedings of the 16th Annual Conference of Animal Science Association of Nigeria.* p.133-137.

Kamboj, V.P. (2000). Herbal Medicine. *Current Science*; 78:35-51.

Kamis, A.B., Modu, S., Zanna, H.,&Oniyangi, T. A.(2003). Preliminary biochemistry and haematological effects of aqueous suspension of pulp of hyphaenethebaica (1) mart in rats. *Biokemistri.* 13 (1): 1-7.

Karaye, I. U., Aliero, A. A., Muhammad, S., & Bilbis, L. S. (2013). Evaluation of Nutrient and Antinutrient Content of selected Nigerian cucurbita seed. Research journal of Pharmaceutical, Biological and chemical sciences. 4(1) 137-142

Katzir N, Danin-Poleg Y, Tzuri G, Karchi Z, Lavi U, Cregan PB (1996). Length polymorphism and homologies of microsatellites in several Cucurbitaceae species. Theor. Appl. Genet. 93: 1282-1290.

Kensler, T. W., Wakabayashi, N., and Biswal, S. (2007). Cell survival responses to environmental stresses via the Keap1-Nrf2-ARE pathway. Annu Rev Pharmacol Toxicol 47, 89–116.

Klaunig, J. E., and Kamendulis, L. M. (2004). The role of oxidative stress in carcinogenesis. Annu Rev Pharmacol Toxicol 44, 239–67.

Krantz SB, Shields MA, Dangi-Garimella S, Cheon EC, Barron MR, Hwang RF, et al. MT1-MMP cooperates with Kras (G12D) to promote pancreatic fibrosis through increased TGF-β signaling. Mol Cancer Res. 2011;9:1294–304.

Katalinic V, Milos M, Kulisic T, Jukic M (2006). Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. *Food Chemistry*, 94: 550–557.

Katsurano, M; Niwa, T; Yasui, Y; Shigematsu, Y; Yamashita, S; Takeshima, H; Lee, M S; Kim, Y-J; Tanaka, T; Ushijima, T (2011). "Early-stage formation of an epigenetic field defect in a mouse colitis model, and non-essential roles of T- and B-cells in DNA methylation induction". *Oncogene*. 31 (3): 342–351. doi:10.1038/onc.2011.241. PMID 21685942

Khan, M. Z., Szarek, J., Koncicki, A., and Krasnodebska-Depta, A. (1994). Oral administration of monensin and lead to broiler chicks: effect on haematological and biochemical parameters. *Acta Veterinaria Hungarica, 42*(1), 111-120.

Khan, T. A., & Zafar, F. (2005).Haematological Study in Response to Varying Doses of Estrogen in Broiler Chicken. *International Journal of Poultry Science, 4*(10), 748-751.

Kilani S, Sghaier M, Limem L, Bouhlel I, Boubaker J Bhouri W, Skandrani L, Neffatti A, Ammarb R, Dijoux- Franca M, Ghedira K, Chekir-Ghedira L (2008). *In vitro* evaluation of antibacterial, antioxidant, cytotoxic and apoptotic activities of the tubers infusion and extracts of *Cyperus rotundus. Bioresource Technology*. 99: 9004-9008.

Kirtikar K. R., Basu B.D. (2009).Indian Medicinal Plants, 2nd edition, vol 9, Oriental enterprises; p. 1607

Kitts D, Yuan Y, Wijewickreme AN, Hu C (2000). Antioxidant properties of a North American ginseng extract. *Molecular Cellular Biochemistry*, 203: 1-10.

Krebs, C. (1928). Experimenteller Alkoholkrebs bei weissen Ma¨usen. Z. Immun. Exp. Therapeutics, 50, 203–218.

Lambert R, Sauvaget C, Sankaranarayanan R. Mass screening for colorectal cancer is not justified in most developing countries. Int J Cancer. 2009; 125(2):253-256.

Larson R.A., The antioxidants of higher plants, Phytochemistry, 1988,27, 969-78.

Lahon, L.C. (1978). Preliminary Pharmacological and Anticestodal Screening Of *Cucurbita Maxima. Indian Journal Pharmacy*; 10:315-317**.** IJPP VOL 2 ISSUE 3

LaMont, J.T. (1978). Experimental colon cancer. *Gastroenterology*, 75, 1157–1169.

Laqueur,G.L. (1964). Carcinogenic effects of cycad meal and cycasin, methylazoxymethanol glycoside, in rats and effects of cycasin in germfree.

Laqueur, G.L. (1963). Carcinogenic Properties of nuts from Cycas circinalis L. indigenous to Guam. *Journal National Cancer Institute*, 31, 919–951.

Lee, S.E., Hyun, J.H., Ha, J., Jeong, H., Kim, J.H. (2003). Screening of medicinal plant extracts for antioxidant activity. *Life Science*, 73: 167–179.

Lee, Y.K., Chung, W.I. and Ezura, H. (2003). Efficient Plant Regeneration via Organogenesis in Winter Squash (Cucurbita maxima Duch.). *Plant Science* 164, 413 -418.

Lim, L.P., Lau, N.C., Garrett-Engele, P., Grimson, A., Schelter, J.M., Castle, J., Bartel, D.P., Linsley, P.S., Johnson, J.M. (2005). "Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs". *Nature*. 433 (7027): 769–773. doi:10.1038/nature03315.PMID 15685193.

Lim, Y. Y., Lim, T. T. and Tee, J. J. (2006). Antioxidant properties of guava fruit: Comparison with some local fruits. *Sunway Academic Journal* 3: 9-20.

Lisco H. (1947). Carcinoma of the colon in rats following the feeding of radioactive yttrium *Cancer Research.* 7; 721-725.

Lisco, H. (1947). Carcinoma of the colon in rats following the feeding of radioactive yttrium. *Cancer Research* 7, 721-725.

Lopez, M., Martinez, F., Del Valle, C., Ferrit, M., Luque, R. (2003). Study of phenolic compounds as natural antioxidants by a fluorescence method. Talanta 60: 609-616.

López, A. P., Gochicoa, M. T. T. and Franco, A. R. (2010). Activities of antioxidant enzymes during strawberry fruit development and ripening. *Biological Plantarum* 54(2): 349-352.

Lopez-Lazaro M (August 2015). "Stem cell division theory of cancer". *Cell Cycle*. 14 (16): 1–2. doi:10.1080/15384101.2015.1062330. PMID 26090957.

Maeda S, Kamata H, Luo J L, Leffert H, Karin M (2005). IKKbeta couples hepatocyte death to cytokinedriven compensatory proliferation that promotes chemical hepatocarcinogenesis. Cell, 121(7):977– 990

Mantovani A, Allavena P, Sica A, Balkwill F (2008). Cancer-related inflammation. Nature, 454(7203): 436–444

Marnett, L. J. (2000). Oxyradicals and DNA damage. Carcinogenesis 21, 361–70.

Martin J.P Jr, Daily M, Sugarman E. (1987). Negative and positive assays of superoxide dismutase based on hematoxylin autoxidation

Masson V, de la Ballina LR, Munaut C, Wielockx B, Jost M, Maillard C, et al. Contribution of host MMP-2 and MMP-9 to promote tumor vascularization and invasion of malignant keratinocytes. FASEB J. 2005;19:234–6.

Miyamoto, hf., and Takizawa, S.: Brief communica- tion-Carcinoma of highly inbred rats. J. Natl. Cancer Inst. 55 :1471-1472, 1975.

Manson, M.M. (2003). Cancer Prevention-the Potential for diet to modulate molecular signaling. *Trends Molecular Medicine* 9; 11-18.

Manzoor, M., Anwar, F., Mahmood, Z., Rashid U. and Ashraf, M. (2012). Variation in minerals, phenolics and antioxidant activity of peel and pulp of different varieties of peach (*Prunus persica* L.) fruit from Pakistan. *Molecules*, 17: 6491.

Maton, A., Hopkins, R. L. J., McLaughlin, C. W., Johnson, S., Warner, C. W., LaHart, D., and Wright, J. D. (1993). *Human Biology and Health*. Englewood Cliffs, New Jersey, USA: Prentice Hall. 63:132-135.

Merck Manual, (2012). Haematologic reference ranges. *Mareck Veterinary Manual*. Retrieved from http://www.merckmanuals.com/. On 23rd, January, 2017.

Mitra, P., Ramaswamy S. H., Chang S.K. (2009). Pumpkin (*Cucurbita maximus*) seed oil extraction using supercritical carbon dioxide and physicochemical properties of the oil. *Journal of Food Engineering*. 95:208-213.

Miwa, M. (1976). Spontaneous colon tumors in rats. *Journal National Cancer Institute*, 56, 615–621.

Mitruka, B. M., and Rawnsley, H. M. (1977). Clinical biochemical and hematological reference values in normal experimental. *USA: Masson Publishing Inc*. (p.134-135).

Miyamoto,M. *et al*., (1991). Adenocarcinoma in the ascending colon of ACI strain rat foster bred by WF-Osaka female rat. *Medline Journal Osaka University*, 40, 1–3

Mmereole, F. U. C. (2008). The Effects of Replacing Groundnut Cake with Rubber Seed Meal on the Haematological and Serological Indices of Broilers. *International Journal of Poultry Science, 7*(6), 622-624.

Nambiar,P.R. (2003). Preliminary analysis of azoxymethane induced colon tumors in inbred mice commonly used as transgenic/knockout progenitors. *International Journal Oncology*, 22, 145–150.

Moore, S. and Stein, W. H. 1948. Photometric ninhydrin method for use in the chromatography of amino acids. Journal of Biological Chemistry 176: 367-388.

Muntean E, Nicoleta M, Marcelmatei D. (2013). University of AgriculturalSciences and Veterinary Medicine Cluj-Napoca, RO. Hop and medicinal plants, Year XXI, No. 1-2 (41-42)

Murphy G, Devesa SS, Cross AJ, Inskip PD, McGlynn KA, Cook MB. Sex disparities in colorectal cancer incidence by anatomic subsite, race and age. Int J Cancer. 2010;128: 1668-1675.

National Center for Health Statistics, Division of Health Interview Statistics. National Health Interview Survey Public Use Data File 2010. Centers for Disease Control and Prevention. Hyattsville, MD., 2011.

Nakane PK, Pierce GB: Enzyme-labelled antibodies: preparation and application for localization of antigen. J Histochem Cytochem 14:929. 1966

National Cancer Institute. DevCan: Probability of Developing or Dying of Cancer Software, Version 6.7.0; Statistical Research and Applications Branch, National Cancer Institute, 2005.

Newman DJ, Cragg GM, Snader KM. Natural products as sources of new drugs over the period 1981-2002. J Nat Prod. 2003;66:1022–37

Neufert, C. et al. (2007). An inducible mouse model of colon carcinogenesis for the analysisof sporadic and inflammation-driven tumor progression. Nat. *Protoc.*, 2, 1998-2004

Newmark, H.L. (2001). A western-style diet induces benign and malignant neoplasms in the colon of normal C57Bl/6 mice. Carcinogenesis,

Nwaoguikpe, R. N, Ujowundu, C.O & Okwu, G. N(2013).The antisickling potentials of four cucurbits (T occidentalis, C maxima, C sativus and C lonatu). Sch. J. App. Med. Sci. 1(3):191-198.

Okazaki I M, Kotani A, Honjo T (2007). Role of AID in tumorigenesis. Adv Immunol, 94: 245–273

Ogunbajo, S. O., Alemede, I. C., Adama, J. Y., and Abdullahi, J. (2009). Haematological parameters of Savannah brown does fed varying dietary levels of flamboyant tree seed meal. *Proceedings of 34th Annual Conference of Nigerian Society for Animal roduction.* p. 88-91.

Okunlola, D. O., Olorunisomo, A. O., Aderinola, A. O., Agboola, A. S., and Omole, O. G. (2012). Haematology and serum quality of red Sokoto goats fed Baobab (Adansonia digitata) fruit meal as supplement to guinea grass *(Panicum maximum). Proceedings of the 17th Annual Conference of Animal Science Association of Nigeria.* p.427-433.

Olafedehan, C. O., Obun, A. M., Yusuf, M. K., Adewumi, O. O., Oladefedehan, A. O., Awofolaji, A. O., and Adeniji, A. A. (2010). Effects of residual cyanide in processed cassava peal meals on haematological and biochemical indices of growing rabbits *Proceedings of 35th Annual Conference of Nigerian Society for Animal Production.* p.212.

Omiyale, C. A., Yisa, A. G., and Ali-Dunkrah, L. A. (2012). Haematological characteristics of Yankasa sheep fed fonio (Digitaria iburua) straw based diets. *Proceedings of 37th Annual Conference of Nigerian Society for Animal Production.* p. 87-89.

Onwuka, G. I. (2005). Food analysis and instrumentation: theory and practice. *Naphathali prints, Nigeria,* 95-96.

Onyeyili, P. A., Egwu, G. O., Jibike, G. I., Pepple, D. J., and Ohaegbulam, J. O. (1992). Seasonal variation in haematological indices in the grey-breasted guinea fowl (*Numida mealagris Gallata pallas*). *Nigerian Journal of Animal Production*, *18*(2), 108-110.

Orwa, C., Mutua, A., Kindt,R., Jamnadass, R., Simons, A. (2009). "Agroforestree database: a tree reference and selection guide version 4.0". Url: http://www.Worldagroforestry.org/af/treedb/ (Accessed on 15 February, 2011).

Ovuru, S. S., and Ekweozor, I. K. E. (2004). Haematological changes associated with crude oil ingestion in experimental rabbits. *African Journal of Biotechnology, 3*(6), 346-348.

Oyawoye, B. M., & Ogunkunle, H. N. (2004). Biochemical and haematological reference values in normal experimental animals. *New York: Masson*. p. 212-218.

Pearson, D. (1976): Chemical Analysis of Foods, 7 th edition, Edinburgh, Churchil Living Stone, pp 204 – 210

Poonia P, Junaid N, Chaudhary G: In-Vitro antioxidant potential of jasminum mensy hance (leaves) extracts. Research journal of pharmaceutical, biological and chemical science 2011; 2(1):348- 357.

Przybylowska K, Kabzinski J, Sygut A, Dziki L, Dziki A, Majsterek I. An association selected polymorphisms of XRCC1, OGG1 and MUTYH gene and the level of efficiency oxidative DNA damage repair with a risk of colorectal cancer. Mutat Res. 2013; 745-746:615.

Peters, S. O., Gunn, H. H., Imumorin, I. G., Agaviezor, B. O., and Ikeobi, C. O. (2011). Haematological studies on frizzled and naked neck genotypes of Nigerian native chickens. *Tropical Animal Health Production, 43*(3), 631-638.

Piao M, Kang k, Dong R, OkKo D, Wang Z, You H, Kim H, Kim J, Kang S, Hyun J (2008). Hyperoside prevents oxidative damage induced by hydrogen peroxide in lung fibroblast cells via an antioxidant effect. *Biochimica Biophysica Acta.* 1780 : 1448–1457.

Poole, T. B. (1987). The UFAW Handbook on the Care and Management of Laboratory and other research Animals. *Universities Foundation for Animal Welfare, Longman Scientific and Technical, Harlow*, U.K. 6th ed.

Popoola. A.O., Omodele, F.O., Oludara, M.A., Ibrahim, N.A., Igwilo, A.I. and Makanjuola, S.B.L. (2013). Prevalence and pattern of cancers among Adults Attending a Tertiary Health Institution in Lagos, Nigeria. *Medical Journal,* 85(4): S4-S6.

Purves, W. K., Sadava, D., Orians, G. H., and Heller, H. C. (2003). Life*: The science of Biology* 7th ed., p.954.

Rahal A, Singh V, Mehra D, Rajesh S, Ahmad A.H (2014). Prophylactic efficacy of *Podophyllum hexandrum* in alleviation of immobilization stress induced oxidative damage in rats. *Journal of Natural Products,* vol. 2, pp. 110-115

Rikardsen OG. Prognostic markers in oral squamous cell carsinoma. Faculty of health science: UiT- The arctic university of Norway; 2014.

Richie,J.P.Jr et al. (2007) Induction of colon tumorigenesis by glutathione depletion in p53-knock- out mice. Int. J. Oncol., 30, 1539–1543.

Robinson, R.W. and Decker-Walters, D. S. 1997. Cucurbits. New York: CAB International.

Robinson, R. w and Decker Walter, D.S (1997), "What are cucurbits", R.W. Robinson and D.S. Decker - Watters, CAB international, New York. Pg 1- 22

Rodríguez D,Morrison C J, Overall CM. Matrix metalloproteinases: what do they not do? New substrates and biological roles identified by murine models and proteomics. Biochim Biophys Acta. 1803;2010:39–54.

Rodriguez-Amaya, D. B. 1999. A guide to carotenoid analysis in foods. Washington: International Life Sciences Institute (ILSI) Press.

Rosenberg, S.A. Karnofsky Memorial Lecture. The immunotherapy and gene therapy of cancer. J. Clin. Oncol. 1992, 10, 180.

Rowlatt,C. et al. (1969) Naturally occurring tumors and other lesions of the digestive tract in untreated C57BL mice. J. Natl Cancer Inst., 43,1353–1364.

Rice-Evans C, Miller N, Paganga G (1996). Structure–antioxidant activity relationships of flavonoids and phenolic acids. *Free Radical Biology Medicine,* 20: 933–956.

Richie, J. P. Jr. (2007). Induction of colon tumorigenesis by glutathione depletion in p53-knock-out mice. *Internet Journal Oncology*, 30, 1539–1543.

Robinson, R.W. and Decker-Walters, D. S. (1997). *Cucurbits.* New York: CAB International.

Rosenberg, D.W. (1995). Induction of aberrant crypts in murine colon with varying sensitivity to colon carcinogenesis. *Cancer Lett.,* 92, 209–214.

Rowlatt, C., Tapsell, L.C., Hemphill, I. and Cobiac, L. (1969). Naturally occurring tumors and other lesions of the digestive tract in untreated C57BLmice. *Journal National Cancer Institute*, 43, 1353–1364. 29.

Saha P, Mazumder UK and Haldar P.K. (2011). Evaluation of hepatoprotective activity of *Cucurbita maxima* aerial parts. *Journal of Herbal and Toxicology*; 5(1):17-22.

Saha P, Mazumder UK, Haldar PK, Bala A, Kar B, Naskar S. (2011). Evaluation of hepatoprotective activity of *Cucurbita maxima* aerial parts. *Journal Herbal Medline Toxicology*; 5(1): 17-22.

Saha P, Mazumder UK, Haldar PK, Naskar S, Kundu S, Bala A, Kar B. (2011). Anticancer activity of methanol extract of Cucurbita maxima against Ehrlich ascites carcinoma. *Internent Journal Research Pharmacy Science*; 2(1): 52-59.

Saha P., Bala A., Kar B., Naskar S., Mazumder U.K., Haldar P.K., Gupta M., (2011), Antidiabetic activity of *Cucurbita maxima* aerial parts. *Research Journal of Medicinal Plant,* 5, 577-586.

Saha P., Mazumder U. K., Haldar P. K., Asis B., Biswakanth K., Sagar N., (2011), Evaluation of hepatoprotective activity of *Cucurbita maxima* aerial parts. *Journal of Herbal Medicine and Toxicology*, 5, p.17-22.

Saha P., Mazumder U.K., Haldar P.K., Naskar S., Kundu S., (2011), Anticancer activity of methanol extract of Cucurbita maxima against Ehrlich as-cites carcinoma. *Internent Journal Research Pharmacological Science*,1, p.52-59.

Saha, P. (2011). Evaluation of Hepatoprotective Activity of *Cucurbita maxima* Aerial Parts. *Journal of Herbal Medicine and Toxicology*; 5:17-22.

Sidell, B. D., and O' Brien, K. M. (2006). When bad things happen to good fish: the loss of haemoglobin and myoglobin expression in Antarctic icefishes. *The Journal of Experimental Biology, 209,* 1791-1802.

Soetan, K. O., Akinrinde, A. S., and Ajibade, T. O. (2013). Preliminary studies on the haematological parameters of cockerels fed raw and processed guinea corn *(Sorghum bicolor)* (p. 49-52). *Proceedings of 38th Annual Conference of Nigerian Society for Animal Production. Some Traditional Herbal Medicines*. 82:43-68.

Sopan, B.A., Vasantrao, D.N. and Ajit, S.B. (2014). Total Phenolic Content and Antioxidant Potential of Cucurbita Maxima (Pumpkin) Powder. *International Journal of Pharmacological Science Research.* 5(5): 1903-07.doi: 10.13040/IJPSR.0975-8232.5 (5).1903-1907.

Souri, E., G. Amin, H. Farsam and T.M. Barazandeh. (2008). Screening of antioxidant activity and phenolic content of 24 medicinal plant extracts. *DARU,* 16: 83-87.

SrijitaDutta (2015). Natural sources as potential anti-cancer agents: A review. Department of Pharmacology, NSHM college of pharmaceutical technology, Kolkata. ISSN: 2348-8948. Vol: 3; Issue: 5.

Sharma and Ramana Rao, T. V. (2013)"Nutritional Quality Characteristics Of Pumpkin Fruit As Revealed By Its Biochemical Analysis"International Food Research Journal (2013)20(5): 2309-2316.

Singh S, Singh H, Singh PP, Murad MH, Limburg PJ. Anti-Diabetic Medications and the Risk of Colorectal Cancer in Patients with Diabetes Mellitus: A Systematic Review and Meta-Analysis. Cancer Epidemiol Biomarkers Prev. 2013;22: 2258-2268

Singh J, Kulkarni N, Kelloff G, Reddy BS. Modulation of azoxymethane-induced mutational activation of ras protooncogenes by chemopreventive agents in colon carcinogenesis. Carcinogenesis 1994;15: 1317.

Singh J, Rivenson A, Tomita M, Shimamura S, Ishibasi N, Reddy BS. Bifidobacterium longum, a lactic acid-producing intestinal bacterium inhibits colon cancer and modulates the intermediate biomarkers of colon carcinogenesis. Carcinogenesis 1997;18:833

Siddhartha, S., A. Mehta and P. Mehta. (2011). Hepatoprotective activity of *Cajanus cajan* against Carbon tetrachloride Induced Liver Damage. *Internet Journal Pharmacy Pharmacological Science*, 3: 0975_1491.

Simpson K.L, Kamata T., (1979), Use of carotenoids in fish feeds. Finfish Nutr.Finfish Technol. 2, p.415-424

Sinha AK.(1972). Colorimetric assay of catalase. Anal Biochem.47: 389

Smith G, Carey FA, Beattie J, Wilkie MJ, Lightfoot TJ, Coxhead J, et al. Mutations in APC, kirsten-ras, and p53--alternative genetic pathways to colorectal cancer. Proc Natl Acad Sci U S A. 2002; 99: 9433-8.

Smith J, Tho LM, Xu N, Gillespie DA. The ATM-Chk2 and ATRChk1 pathways in DNA damage signaling and cancer. Adv Cancer Res. 2010; 108: 73-112.

Smith JSC, Smith OS (1989). The description and assessment of distances between inbred lines of maize: the utility of morphological, biochemical and genetic descriptors and a scheme for the testing of distinctiveness between inbred lines. Maydica 34: 151161.

Smyth M J, Dunn G P, Schreiber R D (2006). Cancer immunosurveillance and immunoediting: the roles of immunity in suppressing tumor development and shaping tumor immunogenicity. Adv Immunol, 90: 1–50

Sopan BA, Vasantrao DN and Ajit SB: Total Phenolic Content and Antioxidant Potential of Cucurbita Maxima (Pumpkin) Powder. Int J Pharm Sci Res 2014; 5(5): 1903-07.doi: 10.13040/IJPSR.0975-8232.5 (5).1903-07

Sturgeon CM, Duffy MJ, Stenman UH, *et al*; National Academy of Clinical. Biochemistry laboratory medicine practice guidelines for use of tumor markers in testicular, prostate, colorectal, breast, and ovarian cancers. Clin Chem. 2008 Dec;54(12):e11-79.doi: 10.1373/clinchem.2008.105601.

Sturgeon CM, Lai LC, Duffy MJ; Serum tumour markers: how to order and interpret them. BMJ. 2009 Sep 22;339:b3527. doi: 10.1136/bmj.b3527

Sukumar S, Notario V, Martin- Zanca D, Barbacid M. (1983). Induction of mammary carcinomas in rats by nitroso-methylurea involves malignant activation of H-ras-1 locus by single point mutation. Nature 306 (5944):658-61

Surveillance, Epidemiology, and End Results (SEER) Program SEER Stat Database: Mortality – All COD, Aggregated With State, Total U.S. (1969-2010) <Katrina/Rita Population Adjustment>, National Cancer Institute, DCCPS, Surveillance Research Program, Cancer Statistics Branch, released April 2013. Underlying mortality data provided by NCHS 2013.

Tariq K, Ghias K. Colorectal cancer carcinogenesis: a review of mechanisms. Cancer Biol Med. 2016; 13: 120-35. doi: 10.28092/j.issn.2095-3941.2015.0103

Thomson HJ and Adlakha H (1991). Dose responsive induction of mammary gland carcinomas by the intraperitoneal injection of 1-methyl-1-nitrosourea. Cancer Res **51:**3411-3415

Thomson M, Ali M (2003). Garlic [*Allium sativum*]: a review of its potential use as an anti-cancer agent. Curr Cancer Drug Targets, 3, 67-81.

Thomson DMP, Krupey J, Freedman SO, Gold P. The radioimmunoassay of circulating carcinoembryonic antigen of the human digestive system. Proc Natl Acad Sci USA 1969;64:161-7.

Thomas P, Toth CA, Saini KS, Jessup JM, Steele G. The structure, metabolism and function of the carcinoembryonic antigen gene family. Biochim Biophys Acta 1990;1032:177-89

Thompson, J. E., Ledge, R. L., Barber, R. F. 1987.The role of free radicals in senescence and wounding. New Phytologist 105: 317-344.

Thompson J, Grunert F, Zimmermann W. carcinoembrayonic antigen gene family: molecular biology and clinical perspectives. J Clin Lab Anal 1991;5:344-66

Thompson, J. E., Ledge, R. L., Barber, R. F. (1987).The role of free radicals in senescence and wounding. *New Phytologist*. 105: 317-344.

Tiwari A.K. (2001). Imbalance in antioxidant defense and human diseases: Multiple approach of natural antioxidant therapy. *Current Science*, 8: 1179–1187.

Togun, V. A., Oseni, B. S. A., Ogundipe, J. A., Arewa, T. R., Hammed, A. A., Ajonijebu, D. C., Mustapha, F. (2007). Effects of chronic lead administration on the haematological parameters of rabbits – a preliminary study. *Proceedings of the 41st Conferences of the Agricultural Society of Nigeria*. p. 341.

Toth,B., Subash, V. K., Saritha, G. (1976). Production of intestinal and other tumours by 1,2-dimethylhydrazine dihydrochloride in mice. II. Scanning electron microscopic and cytochemical study of colonic neoplasms. Br. *Journal Express Pathology*, 57, 696–705.

Ugwuene, M. C. (2011). Effect of Dietary Palm Kernel Meal for Maize on the Haematological and Serum Chemistry of Broiler Turkey. *Nigerian Journal of Animal Science, 13*, 93-103.

Valko, M., Rhodes, C. J., Moncol, J., Izakovic, M., and Mazur, M. (2006). Free radicals, metals and antioxidants in oxidative stress-induced cancer. Chem Biol Interact 160, 1–40.

Vivek, K.G. and Surendra, K.S.: Plants as natural antioxidants. Natural. Production. Radia., 5: 326-334 (2006).

Von Sonntag, C. (1987). New aspects in the free-radical chemistry of pyrimidine nucleobases. Free Radic Res Commun 2, 217–24.

Vandenberghe,J. (1985). Spontaneous adenocarcinoma of the ascending colon inWistar rats: the intracytoplasmic presence of a Campylobacterlike bacterium. *Journal Comp. Pathology*, 95, 45–55.

Verma, S. (2008). Current and future status of herbal medicines. *Veterinary World*; 1:347-350.

Villasefior,M.I., Lemon, P. Antigenotoxic spinasterol from *Cucurbita maxima* flowers. *Mutation Research* 1996; 360: 89-93.

Vogelstein, Bert; Kinzler, Kenneth W. (2004). "Cancer genes and the pathways they control". *Nature Medicine*.**10** (8): 789–99. doi:10.1038/nm1087. PMID 15286780.

Walpole,A.L. (1952) The carcinogenic action of 4-aminodiphenyl and 3:2#-dimethyl-4-amino-diphenyl. Br. *Journal Indian Medicine* 9, 255–263.

Wang W., Xia M.X., Chen J., Yuan R., Deng F.N. and Shen F.F. (2016). Gene expression characteristics and regulation mechanisms of superoxide dismutase and its physiological roles in plants under stress. Biochemistry (Mosc.) 81 (5): 465–480.

Weisiger, A. A., and Fridovich, I. Mitochondrial superoxide dismutase. Site of synthesis and intramitochondrial localization. J. Biol. Chem., 248: 4793-4796, 1973

Weisburger JH. Causes of gastric and esophageal cancer: Possible approach to prevention by vitamin C. Int J Vit Nutr Res 1985; 27:38 1- 402

Weisburger, J. H., Fiala, E., Narisawa, T., and Reddy, B. S. : On the etiology and prevention of cancer in the gastrointestinal tract. In Fundamentals in Cancer Pre- vention, P. N. Magee et al., Eds. Baltimore, University Park Press, 1976; pp. 121-142.

Ward, J.M. (1974). Morphogenesis of chemically induced neoplasms of the colon and small intestine in rats. *Laboratory Invest*., 30, 505–513.

Weisburger, J. H., Reddy, B. S., and Joftes, D. L.: Colo-Rectal Cancer. Geneva, UICC Technical Report Series, Vol. 19, 1975.

Westermarck J, Kähäri V.M. Regulation of matrix metalloproteinase expression in tumor invasion. FASEB J. 1999;13:781–92.

Weisenberger DJ, Siegmund KD, Campan M, Young J, Long TI, Faasse MA, et al. CpG island methylator phenotype underlies sporadic microsatellite instability and is tightly associated with BRAF mutation in colorectal cancer. Nat Genet. 2010; 38: 787-93.

Welsch CW. Host factors affecting the growth of carcinogen-induced rat mammary carcinomas: A review and tribute to Charles Brenton Huggins. Cancer Res 1985; 45:3415-3443.

Whitaker, T. W. and Davis, G. N. 1962. Cucurbits. New York: Interscience Publishers

William CS, Kwok N L (2007). In vitro and in vivo anti-tumor effects of Astragalus membranaceus. Cancer Letters, 252, 43-54. Woo HY, Park KY, Rhu CH, et al (2006). β-Lapachone, a Quinone Isolated from Tabebuia avellanedae, Induces apoptosis in HepG2 hepatoma cell line through induction of bax and activation of caspase. J Med Food, 9, 161-8.

Wilson CL, Heppner KJ, Labosky PA, Hogan BL, Matrisian LM. Intestinal tumorigenesis is suppressed in mice lacking the metalloproteinase matrilysin. Proc Natl Acad Sci U S A. 1997;94:1402–7.

Waugh, A., Grant, A. W., & Ross, J. S. (2001). Ross and Wilson Anatomy and Physiology in Health and Illness. *Churchill Livingston, an imprint of Elsevier Science Limited*. 9th edition, p. 59-71.

Weisburger,J.H. et al. (1977). Colon cancer: its epidemiology and experimental production. *Cancer,* 40, 2414–2420.

Whitaker T.W., Bemis W.P., (1975). The origin and evolution of the cultivated cucurbitae. *Bull.Tom. Bot.Club*, 102, p.362-368.

Whitaker, T. W. and Davis, G. N. (1962). *Cucurbits*. New York: Interscience Publishers.

Woo; Lee, Jae-Hyuk (2011). "Promoter methylation status of hMLH1, hMSH2, and MGMT genes in colorectal.

Wood D, Jenne J. (2007). The conservation of agrobiodiversity on-farm: Questioning emerging paradigm. *Biodiversity. Conservation*. 6: 109-129.

Woods MO, Williams P, Careen A, Edwards L, Bartlett S, Mclaughlin JR, et al. A new variant database for mismatch repair genes associated with Lynch syndrome. Hum Mutat. 2007; 28: 669-73.

Wood, L.D. *et al*. (2007) The genomic landscapes of human breast and colorectal cancers. Science, 318, 1108–1113.

World Health Organization. Global Status Report on Non Communicable Disease; 2010. Accessed 06 June 2011. Available: http://www.who.int/chp/ncd_global_status_report/en/.

Wu T, Zhou J, Zhang Y, Cao J (2007). Characterization and inheritance of a bush-type in tropical pumpkin (Cucurbita moschata Duchesne). Scientia Horticulturae, pp. 1-4.

Yao, J.K., Ravinder, D.R. and Daniel, P. (1999). Human Plasma Glutathione

Yeatman, T.J. (2001). Colon Cancer. Encyclopedia of Life Sciences. Macmillan Publishers Ltd, Nature Publishing Group.

Zia-Ul-Haq, M., S. Ahmad, L. Calani, T. Mazzeo, D.D. Rio, N. Pellegrini and V. DeFeo, 2012a. Compositional study and antioxidant potential of *Ipomoea hederacea* Jacq. and *Lepidium sativum* L. seeds. *Journal of Molecular Science,*, 17: 10306-10321.

Zia-Ul-Haq, M., S.A. Shahid, S. Ahmad, M. Qayum and I. Khan, 2012b. Antioxidant potential of various parts of *Ferula assafoetida* L. *Journal on Medicinal Plant Research,* 6: 3254-3258.

Zubair, M., F. Anwar and S.A. Shahid, 2012. Effect of extraction solvents on phenolics and antioxidant activity of selected varieties of Pakistani rice (*Oryza sativa* L.). *International Journal Agriculture Biology,* 14: 935-940.

Zhou Y, Yu C, Miao X, Tan W, Liang G, Xiong P, *et al.* Substantial reduction in risk of breast cancer associated with genetic polymorphisms in the promoters of the matrix metalloproteinase-2andtissue inhibitor of metalloproteinase-2 genes. Carcinogenesis.

2004;25: 399–404Xu E, Xia X, Lü B, Xing X, Huang Q, Ma Y, et al. Association of matrix metalloproteinase-2 and -9 promoter polymorphisms with colorectal cancer in Chinese. Mol Carcinog. 2007;46:924–9.

APPENDICES 1.0

Effect of *Cucurbita maxima* fruit on Percentage Packed Cell Volume (PCV), Neutrophil and Lymphocyte following 12 Weeks MNU Administration in the Blood of Different Experimental Group.

Group	Treatment	PCV(%)	NEUTRO(%)	LYMPHO(%)

i	**Normal Control**	**42.30±2.2^{b}**	**26.30±6.2^{ab}**	**52.70±4.5^{b}**
ii	**MNU Control**	**32.20±4.2^{a}**	**31.80±5.3^{c}**	**66.00±3.5^{c}**
iii	**MNU+2.5% of *C. maxima* fruit**	**36.20±2.4^{a}**	**29.30±2.4^{b}**	**60.30±2.4^{bc}**
iv	**MNU+5% of *C. maxima* fruit**	**40.10±3.1^{b}**	**27.40±3.4^{b}**	**53.50±3.5^{b}**
v	**MNU+10% of *C. maxima* fruit**	**48.80±2.3^{c}**	**24.50±3.1^{a}**	**48.20±1.2^{a}**
vi	**10% of *C. maxima* fruit Control**	**50.40±1.3^{c}**	**22.30±2.5^{a}**	**46.20±3.2^{a}**

Values are Mean±SD; Values having different letters across the column are significantly different ($p \leq 0.05$).

APPENDICES 2.0

Effect of *Cucurbita maxima* Fruit on Haemoglobin and Platelet Concentration in the Blood Serum following 12 Weeks MNU Administration in Different Experimental Group.

Group	Treatment	Hb(g/dl)	TP (g/dl)

i	**Normal Control**	**17.56±1.5^{b}**	**7.70±0.5^{a}**
ii	**MNU Control**	**10.40±1.3^{a}**	**5.90±0.8^{a}**
iii	**MNU+2.5% of *C. maxima* fruit**	**12.10±1.2^{a}**	**7.10±0.2^{a}**
iv	**MNU+5% of *C. maxima* fruit**	**14.50±1.2^{ab}**	**8.20±1.0^{ab}**
v	**MNU+10% of *C. maxima* fruit**	**17.60±0.5^{b}**	**9.90±0.6^{b}**
vi	**10% of *C. maxima* fruit Control**	**19.80±0.4^{c}**	**10.30±0.7^{b}**

Values are Mean±SD; Values having different letters across the column are significantly different ($p \leq 0.05$).

APPENDICES 3.0


Effect of *Cucurbita maxima* Fruit on White Blood Cell and Red Blood Cell in the Blood following 12 Weeks MNU Administration in Different Experimental Group.

Group Treatment	WBC (10^{9}/l)	RBC (10^{12}/l)
i Normal Control	**10.70±2.5^{a}**	**8.80±0.3^{ab}**
ii MNU Control	**12.50±2.1^{c}**	**6.10±0.5^{a}**
iii MNU+2.5% of *C. maxima* fruit	**10.90±3.5^{b}**	**7.30±1.0^{a}**
iv MNU+5% of *C. maxima* fruit	**10.20±3.2^{b}**	**8.10±0.6^{ab}**

v MNU+10% of *C. maxima* fruit	**9.40±0.6^{b}**	**8.93±0.5^{ab}**
iv 10% of *C. maxima* fruit Control	**7.90±1.3^{a}**	**9.20±0.8^{b}**


Values are Mean±SD; Values having different letters across the column are significantly different ($p \leq 0.05$).

APPENDICES 4.0




Effect of Dietary *Cucurbita maxima* Fruit Inclusion on MDA Level following 12 Weeks MNU Administration in Rats. Values are Mean ±SD

APPENDICES 5.0



Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Catalase Activity following 12 Weeks MNU Administration in Rats. Values are Mean ±SD

APPENDICES 6.0



Effect of Dietary *Cucurbita maxima* Fruit Inclusion on Superoxide dismutase Activity following 12 Weeks MNU Administration in Rats.

APPENDICES 7.0

Carcinoembryonic Antigen Concentration of the Serum following 12 Weeks Intrarectal Administration of MNU in Rats Fed with Different Level of Dietary Inclusion of *C. Maxima* Fruit.

Group	Treatment	CEA Concentration (ng/ml)
i	NORMAL Control	122.50±21.00[a]
ii	MNU Control	162.77± 17.65[a]
iii	MNU+2.5% of *C. maxima* fruit	133.06±16.26[a]
iv	MNU+5% of *C. maxima* fruit	119.56±8.37[a]
v	MNU+10% of *C. maxima* fruit	108.46±19.58[a]
vi	10% of *C. maxima* fruit Control	107.97±20.96[a]

Values are Mean±SD; Values having different letters across the column are Significantly different ($p \leq 0.05$).